

# Atlas of Missouri Amphibians and Reptiles for 2021



Richard E. Daniel and Brian S. Edmond



# Atlas of Missouri Amphibians and Reptiles for 2021

Richard E. Daniel  
Division of Biological Sciences  
University of Missouri-Columbia  
114 LeFevre Hall  
Columbia MO 65211  
danielr@missouri.edu

Brian S. Edmond  
Computer Services  
Missouri State University  
901 South National Ave  
Springfield MO 65897  
BrianEdmond@missouristate.edu

Recommended citation:

Daniel, R.E. and B.S. Edmond. 2022.  
Atlas of Missouri Amphibians and Reptiles for 2021.  
<<http://atlas.moherp.org/pubs/atlas21.pdf>>

Updated: 03 August 2022 19:13  
Copyright (C) 1997-2022. All rights reserved.

## Table of Contents

Introduction . . . . .	3
Nomenclature and Taxonomy . . . . .	3
Amphibians and Reptiles of Possible Occurrence . . . . .	4
Erroneously Reported and Non-Native Species . . . . .	5
County Records . . . . .	5
Distributions . . . . .	5
Terrestrial Ecoregions . . . . .	6
Aquatic Subregions . . . . .	7
Natural Divisions and Sections Map . . . . .	8
Level I Terrestrial Ecoregions Map . . . . .	9
Level II Terrestrial Ecoregions Map . . . . .	10
Level III Terrestrial Ecoregions Map . . . . .	11
Level IV Terrestrial Ecoregions Map . . . . .	12
Rivers and Major Streams Map . . . . .	13
Watershed Boundaries Map . . . . .	14
Ecological Drainage Units Map . . . . .	15
County and Surrounding States Map . . . . .	16
Species Locality Maps . . . . .	17
County Records Maps . . . . .	47
Acknowledgements . . . . .	77
Literature Cited . . . . .	77
Appendix A: Sources . . . . .	83



## Introduction

Since 1988, members of the Missouri Herpetological Association have compiled new county distribution records for amphibians and reptiles native to the state (Johnson and Powell 1988; Powell *et al.* 1989, 1990, 1991, 1992, 1993a, 1994, 1995, 1996, 1997; Daniel *et al.* 1998, 1999; Daniel and Edmond 2000, 2001; Daniel *et al.* 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021). Cumulative accounts of these new records were presented in Powell *et al.* (1993b), Powell and Daniel (1997), and Daniel and Edmond (2002b). Johnson (2000) presented revised distribution maps based on a combination of these cumulative accounts and more recent annual compilations. The publication of Johnson (1987, 2000) renewed interest in expanding our knowledge of the distribution of the state's herpetofauna. In order to provide herpetologists working in Missouri with current information that is easily taken into the field, we are presenting revised species distribution and county records maps in a format that can be easily updated as new records are documented.

In 1997, with the initiation of the Missouri Herpetological Atlas Project (MOHAP), a database was established to serve as the basis for verifying new distribution records and tracking changes in individual species distributions within the state (Edmond and Daniel 2022). Qualification for inclusion in the database generally requires a catalogued voucher specimen housed in an institutional collection. However, in order to be valuable in tracking changes in the distribution of individual species, it is important to document historical records. Many of the specimens collected prior to 1960, most notably those reported by Hurter (1911) and Anderson (1965), were documented by specimens that have since been lost or destroyed. In order to provide the most accurate picture of the distribution of the state's herpetofauna we have included literature records that we believe to be valid.

Some records were not included because the identification could not be determined with certainty. Within Missouri, the *Hyla versicolor* complex consists of two broadly sympatric and morphologically indistinguishable species (*H. chrysoscelis* and *H. versicolor*). Specimens collected prior to the separation of the two species or those for which the species identification was not determined by some non-morphological trait were not included.

Currently, the MOHAP database contains 38,079 entries and 33,941 valid, non-duplicated collections. This represents specimens housed in 37 museum collections and cited from 33 historical literature sources; 5,318 documented county records; 10,453 unique localities; and 17,768 unique species / locality combinations. This new total constitutes 271 new database entries since the 2019 edition of the *Atlas* (Daniel and Edmond 2020). The total number of species contained in the current edition of the atlas is 118, comprised of 116 native species and 2 non-native species.

Previous editions of the atlas (Daniel and Edmond 2002a, 2003, 2004, 2005, 2006, 2008, 2010, 2012, 2013, 2014, 2016, 2017, 2018, 2019, 2020) are available upon

request. Readers may contribute new records and read more about project details online (Edmond and Daniel 2022). Questions, comments, and suggestions should be directed to the senior author.

## Nomenclature and Taxonomy

Scientific and common names used in this publication follow Crother *et al.* (2017). The chosen common name represents the most restrictive name available and refers to the subspecies found within the state of Missouri. If two or more subspecies are recognized in the state, the species common name was used. Refer to Appendix B for a complete list of scientific names and common names. Since the publication of Johnson (2000), several systematic studies have necessitated nomenclatural changes of some Missouri species.

Frost *et al.* (2006) concluded that several geographically widespread genera represented polyphyletic groups. Their work necessitated the splitting of several familiar genera and applying new names to the species found in Missouri. Missouri toads formerly placed in the genus *Bufo* have been reassigned to the genus *Anaxyrus* and true frogs of the genus *Rana* are now placed in the genus *Lithobates*.

Two other formerly cosmopolitan genera, *Eumeces* and *Elaphe*, have also undergone revision. Brandley *et al.* (2005) resurrected the genus *Plestiodon* for the clade containing the North American skinks and Utiger *et al.* (2002) placed the North American ratsnakes of the genus *Elaphe* into the resurrected genus *Pantherophis*.

A study of the phylogeny of the racers restricted the genus *Coluber* to the New World and also included the coachwhip and whipsnakes of the genus *Masticophis* (Nagy *et al.* 2004). Reeder *et al.* (2002) concluded from their phylogenetic study that the genus *Cnemidophorus* was not a monophyletic assemblage. They presented evidence that species in North America represent a distinct clade and reassigned all U.S. species to the resurrected genus *Aspidoscelis*.

Several studies at the species level have also required changes in the nomenclature of some Missouri reptiles and amphibians. Leaché and Reeder (2002) restricted *Sceloporus undulatus* to the Eastern United States. Missouri populations formerly considered *S. u. hyacinthinus* are now placed in *S. consobrinus*. Starkey *et al.* (2003) determined that southern painted turtles represented a distinct genetic lineage and elevated *Chrysemys dorsalis* to a full species.

Using mitochondrial DNA in a range-wide examination of the *Lampropeltis getula* complex, Pyron and Burbrink (2009) recovered five lineages that they recognized as distinct species. The central lineage, found west of the Mississippi River, was assigned the name of the subspecies widely known in Missouri (speckled kingsnake) and is now known as *L. holbrooki*. More recently, we discovered two black kingsnakes (*L. nigra*) in southeast Missouri (Edmond and Daniel 2014). Because of dramatic shifts in the Mississippi River channel during the Pleistocene, a significant portion of southeast Missouri (*i.e.*, Crowley's Ridge and east) was previously found on the

eastern side of the river. Anderson considered at least some specimens in southeast Missouri as hybrids with the speckled kingsnake (Anderson 1965). Thus, the kingsnakes found in that part of the state are likely black kingsnakes or hybrids with *L. holbrooki*.

Gamble *et al.* (2007) provided molecular evidence that cricket frogs roughly north and west of the Ohio / Mississippi River valleys, including all of Missouri, should be considered a distinct species, *Acris blanchardi*. The common name Blanchard's Cricket Frog is adopted for this species.

Lemmon *et al.* (2007) examined the distributions of North American trilling chorus frogs based on genetics and determined that Missouri populations formerly assigned to the western chorus frog, *Pseudacris triseriata*, were actually the boreal chorus frog, *P. maculata*, and confirmed that the upland chorus frog, *P. feriarum*, of the Mississippi Lowlands was a valid species. In addition, a new species of chorus frog, *P. fouquettei*, has been identified from the interior highlands and western coastal plain (Lemmon *et al.* 2008). Currently, this species is known from three localities in Missouri. Nearby Ozark Plateau localities have been tentatively assigned to *P. feriarum* but are likely to be reassigned to this new species pending further investigation.

Bonett and Chippindale (2004) examined the relationship among members of the *Eurycea multiplicata* complex. They corroborated the conclusion of Thornhill (1990) that Missouri populations assigned to *E. m. griseogaster* were conspecific with *E. tynerensis*. Further, they found that the genus *Typhlotriton* did not show sufficient differentiation from sister taxon *E. tynerensis* to justify recognition and recommended synonymizing it with *Eurycea*.

Crother *et al.* (2011) examined foxsnakes throughout their range and determined that the previously recognized species is composed of eastern and western haplotypes, historically separated by a combination of the Mississippi River and past glaciation events. While it is obvious that foxsnakes in northwest Missouri can be assigned to the western form, the situation in the eastern part of the state is less clear. At least one specimen from southeast Iowa, near the Missouri border, was assigned to the eastern form. Despite their assertion that the Mississippi River formed a barrier, the authors also included the Saint Louis region in their range map for the eastern species, although no specimens from that area were included in their analysis. Thus, we tentatively treat all populations in eastern Missouri along the Mississippi River as the eastern foxsnake (*Pantherophis vulpinus*) and all populations in western Missouri along the Missouri River as the newly described western foxsnake (*P. ramspotti*). Further study of this species complex within Missouri is needed.

Massasaugas in Missouri have persisted only in isolated populations since historical times. Anderson (1965) assigned eastern Missouri populations to *Sistrurus catenatus catenatus*, western populations to *Sistrurus catenatus tergeminus*, and considered animals in the north central part of the state to be hybrids. Recent genetic evidence suggests this taxon actually consists of two distinct species (Kubatko *et al.* 2011). All extant Missouri

populations, including Anderson's intergrades, are readily assigned to the western massasauga (*Sistrurus tergeminus*) (Gibbs *et al.* 2011). Unfortunately, animals from eastern Missouri populations have not been found in more than 75 years, making genetic analysis impossible. We elect to follow Anderson and assign these likely extirpated populations to the eastern massasauga (*Sistrurus catenatus*).

Finally, Shulse (2006, 2007) discovered and described a new locality for Kirtland's Snake (*Clonophis kirtlandii*) in northeast Missouri. In 2007, another adult snake was found dead on a road in Marion County (Daniel 2007). Previously, this species was known in the state from a single Marion County record (Jones 1967). The lack of additional specimens resulted in Johnson (1987, 2000) listing this species as questionable or of possible occurrence in Missouri. The three recent discoveries and the original historical record are included here as an addition to the state's herpetofauna.

## Amphibians and Reptiles of Possible Occurrence

A number of species found in surrounding states are known to occur in close proximity to the borders of Missouri. While not currently recognized as part of the Missouri herpetofauna, some or all of these species may eventually be found within the state.

A number of primarily eastern species have distributions that reach the Mississippi River along a portion of the Missouri border. These include: Southern Two-lined Salamander (*Eurycea cirrigera*), Three-lined Salamander (*Eurycea guttolineata*), Northern Zigzag Salamander (*Plethodon dorsalis*), Northern Slimy Salamander (*Plethodon glutinosus*), Eastern Cricket Frog (*Acris crepitans*), Western Chorus Frog (*Pseudacris triseriata*), Bird-voiced Treefrog (*Hyla avivoca*), Eastern Fence Lizard (*Sceloporus undulatus*), Gray Ratsnake (*Pantherophis spiloides*), and Eastern Ribbonsnake (*Thamnophis sauritus*).

Rivers, even large ones, change course over time and typically form an imperfect geographic barrier. Shepard and Kuhns (2017) examined the separation between Northern Slimy Salamanders (*Plethodon glutinosus*) and Western Slimy Salamanders (*Plethodon albagula*) along the Illinois-Missouri border. They found that salamanders on Fountain Bluff, an island formed of Missouri uplands cut off by post-glacial changes in the Mississippi River channel, belong to the species found in Illinois.

Spotted Dusky Salamander (*Desmognathus conanti*) and Midwestern Earthsnake (*Carphophis amoenus*) have been reported in the southern portion of Crowley's Ridge of Arkansas. Crowley's Ridge is a thin, sandy upland formed during the Pleistocene, which extends from northeastern Arkansas through southeast Missouri to the Shawnee Hills of Illinois. Populations of one or both of these species may exist in the isolated remnants of Crowley's Ridge in Missouri.

Powell *et al.* (2016) shows the distribution of the Western Milksnake (*Lampropeltis gentilis*) extending into

western Missouri along the Kansas border. However, the sparsity of records from this region makes it difficult to accurately define the species boundary. Until additional material is available for evaluation, we have not included this species as part of the state herpetofauna.

## Erroneously Reported and Non-Native Species

Anderson (1945) reported two specimens of the Dwarf Salamander (*Eurycea quadridigitata*) from Roaring River State Park in Barry County. With the nearest known naturally occurring populations of *E. quadridigitata* in southern Arkansas, more than 125 miles from the Missouri border, it is likely that this record is based on misidentified, but morphologically similar Oklahoma Salamanders (*E. tynerensis*).

Johnson and Bader (1974) included Lesser Earless Lizard (*Holbrookia maculata*) based on two specimens collected from Knob Noster State Park in Johnson County. Nickerson and Krager (1972) considered this record problematic because of the close proximity of the park to Whiteman Air Force Base and Central Missouri State University. The nearest record of this species is more than 115 miles away in central Kansas. Further searches failed to produce additional specimens. As a result, this species is not considered to be part of Missouri's herpetofauna.

Anderson (1957) listed the occurrence of the Queensnake (*Regina septemvittata*) in Missouri based on three specimens deposited in the American Museum of Natural History collection. These specimens were ascribed to G.K. Noble and reported to come from Stone County. In further investigation by Roger Conant (1960), Byron C. Marshall, who worked with Noble in the Interior Highlands during this period, reportedly had no recollection of finding this species. The nearest known populations of Queen Snake are in the Boston Mountains of Arkansas, approximately 80 miles south of this locality. The questions about the origin of the specimens and the absence of additional records from Missouri suggest that this species does not occur in the state.

One species of non-native frog and two species of lizards have been reported in Missouri. A population of Greenhouse Frogs (*Eleutherodactylus planirostris*) was established inside a commercial greenhouse in Jefferson City, Missouri (J. Briggler, pers. comm.). The population of these small terrestrial breeding frogs persisted for several years, but has apparently disappeared following renovation of the greenhouse. There are no other reports of this tropical species occurring within the state. Its apparent inability to survive extreme winter weather makes it unlikely for these frogs to exist in Missouri as a free-ranging species outside of sheltered habitats, such as greenhouses.

Mediterranean Gecko (*Hemidactylus turcicus*) is native to the Mediterranean basin and western Asia. This species is easily transported and a highly successful colonizer. Populations of this species have been established worldwide. In the United States, this species is currently established widely throughout the south, especially along

the Gulf and East Coasts. Isolated populations are also found around numerous urban centers of the Midwest and central plains (Powell *et al.* 2016). Two populations appear to be established in Missouri. Bufalino (2004) reported this species from St. Louis County. A second population was discovered in Joplin, Missouri by Mr. A. Braun (Daniel *et al.* 2015).

Briggler *et al.* (2015) reported the presence of a robust population of Italian Wall Lizard (*Podarcis siculus*) in a Joplin neighborhood after being contacted by a resident in 2013. This population apparently descended from a series of lizards imported from Topeka, KS, which escaped in 2001. In many states, including Missouri, it is illegal to release non-native species. Despite prohibitions, the majority of introduced populations of this species in North America originated from released animals in the pet trade (Burke and Deichsel 2008).

Thus far, non-native amphibians and reptiles in Missouri have been confined to urban centers and do not appear to constitute a threat to our native herpetofauna. However, this is not the case everywhere and non-native species that have become invasive are considered by many biologists to be a major threat, second only to habitat loss or degradation, to native species. Responsible pet owners should always exercise care to prevent the spread of this and all non-native species into the natural environment.

## County Records

Earlier versions of this publication used Johnson (2000) as an authority on herpetological county records. However, Johnson's distribution maps are nearly two decades out of date. Furthermore, a number of county records reported by Johnson could not be subsequently verified with museum voucher specimens and are not recognized as valid for the purposes of this project. Beginning with the 2012 version of the *Atlas*, we used only vouchered specimens as the basis for reporting new Missouri county records (except for the aforementioned published records considered valid). Thus, all county records mapped as "open circle" county records in some previous versions of the *Atlas* will not be considered valid and will therefore not be included on the county records maps.

## Distributions

Distribution is one of many characteristics used to describe a species and range maps (such as those found in Conant and Collins (1998)) are used to approximate a species' likely distribution. This atlas uses locality dot maps, with each dot representing a known locality for that species. This conservative method results in an underestimate of a species distribution but is more accurate than a range map. The purpose of the atlas project is to document as closely as possible both current and historical distributions for all native amphibians and reptiles in Missouri.

However, the printed version of the atlas only reflects the best historical approximation of each species'

distribution in the state. The most noticeable resulting incongruity is that fact that some species are shown in historical localities in which they are almost certainly no longer found. For example, the Smooth Greensnake (*Opheodrys vernalis*) was once found in scattered populations in the prairie regions of the state. Due to extensive habitat loss and possibly other reasons, they are extremely rare or extirpated in the state.

While natural communities have long been a staple in ecology, formal classification of natural communities has been undertaken relatively recently in Missouri (Nelson 1987, 2005). Like individual species, natural communities can be described and characterized with distinct distributions and abundances in different ecoregions (see next section). For example, glades are common in the Ozarks but rare elsewhere in the state. Steyermark (1963), Yatskievych (1999), Pflieger (1989), Hawker (1992), and Unklesbay and Vineyard (1992) all describe and summarize the complex interplay among geology, natural history, ecoregions, natural divisions, and natural communities from slightly different perspectives. Taken together, these ideas form the basis of the biogeography of Missouri and they are essential in understanding why plants and animals have the distributions that we see today.

## Terrestrial Ecoregions

An ecoregion classification system attempts to define and describe geographic regions that correspond to broad ecosystem patterns, topography, geology, soils, vegetation patterns, and the distributions of plants and animals. Omernik (1987) described ecoregions (Levels I, II, and III) for the conterminous United States as a hierarchical scheme with Level I corresponding to large regions and Level III representing smaller, more precisely described regions. The Environmental Protection Agency (2003) is coordinating an effort to further subdivide Level III regions into Level IV regions. Nigh and Schroeder (2002) published Level III and Level IV ecoregions for Missouri.

Some amphibian and reptile species follow defined ecoregions closely. For example, the Cave Salamander (*Eurycea lucifuga*) is neatly confined to the Ozark Highlands (Level III) and found throughout the ecoregion, with the exception of most of the Springfield Plateau (Level IV). Several species found in the southeastern alluvial plain are particularly characteristic and also confined to that area (e.g., Three-toed Amphiuma (*Amphiuma tridactylum*) and Southern Watersnake (*Nerodia fasciata*)).

Many species, however, seem to be abundant throughout Missouri, regardless of the region or natural community. The American Bullfrog (*Lithobates catesbeianus*), Snapping Turtle (*Chelydra serpentina*), and Western Ratsnake (*Pantherophis obsoletus*) among several others fall into this category. Finally, many species are more closely associated with a particular natural community or habitat than with a particular ecoregion or natural division. For example, The Flat-headed Snake (*Tantilla gracilis*) can be found in the Ozark Highlands, Osage Prairie, and Interior River Valleys and Hills but it is restricted to rocky glades.

In general, Level IV ecoregions are too granular to be useful in describing the distributions of Missouri amphibians and reptiles, several Ozark salamander species and many of the coastal plains species being notable exceptions. However, this level of detail is useful to understand natural community distributions on which some species are dependent. It also underscores the importance of looking at border states and across artificial political boundaries to understand distributions within Missouri.

Level I (page 8), Level II (page 9), Level III (page 10), and Level IV (page 11) ecoregion maps for Missouri and surrounding states, based on Omernik (1987) and Nigh and Schroeder (2002) are included here. The hierarchy for Levels I, II, and III is included in Table 1 (below), while the Level III and Level IV hierarchy is included in Table 2 (below). Unfortunately, the authors' coding scheme for Level III is inconsistent, though the names do match. Both authors' codes for Level III are included in the tables as a cross-reference. An earlier approach by Thom and Wilson (1980) divided Missouri into natural divisions and sections (page 7), roughly corresponding to Level III and Level IV ecoregions, respectively.

**Table 1.** List of Level I, II, and III Terrestrial Ecoregions from Omernik (1987). Those regions marked with an asterisk (\*) do not occur in Missouri but are sufficiently close to be of interest to Missouri biologists. Level III codes from Nigh and Schroeder (2002) are in parentheses.

- 8. Eastern Temperate Forests
  - 8.3. Southeastern USA Plains
    - 8.3.3. Interior Plateau\* (71)
    - 8.3.6. Mississippi Valley Loess Plains (74)
    - 8.3.2. Interior River Valleys and Hills (72)
  - 8.4. Ozark / Ouachita / Appalachian Forests
    - 8.4.5. Ozark Highlands (39)
    - 8.4.6. Boston Mountains\* (38)
  - 8.5. Southeast US Coastal Plain
    - 8.5.2. Mississippi Alluvial Plain (73)
- 9. Great Plains
  - 9.2. Temperate Prairies
    - 9.2.3. Western Corn Belt Plains (47)
    - 9.2.4. Central Irregular Plains (40)

**Table 2.** List of Level III and Level IV Terrestrial Ecoregions from Nigh and Schroeder (2002). Those regions marked with an asterisk (\*) do not occur in Missouri but are sufficiently close to be of interest to Missouri biologists. Those regions marked with a caret (^) are contained entirely within the state's borders.

- 38. Boston Mountains\* (8.4.6)
  - 38a. Upper Boston Mountains\*
  - 38b. Lower Boston Mountains\*
- 39. Ozark Highlands (8.4.5)
  - 39a. Springfield Plateau
  - 39b. Elk River Hills
  - 39c. White River Hills
  - 39d. Central Plateau
  - 39e. Osage / Gasconade River Hills^
  - 39f. Saint Francois Knobs / Basins^

- 39g. Meramec River Hills^
- 39h. Current River Hills^
- 39i. Easter Ozark Border^
- 39j. Black River Hills Border^
- 39k. Prairie Ozark Border^
- 40. Central Irregular Plains (9.2.4)
  - 40a. Loess Flats and Till Plains
  - 40b. Osage Cuestas\*
  - 40c. Wooded Osage Plains
  - 40d. Cherokee Plains
  - 40e. Claypan Prairie^
- 47. Western Corn Belt Plains (9.2.3)
  - 47d. Missouri Alluvial Plain
  - 47e. Steeply Rolling Loess Prairies
  - 47f. Rolling Loess Prairies
  - 47h. Nebraska / Kansas Loess Hills\*
  - 47i. Loess and Glacial Drift Hills\*
  - 47m. Western Loess Hills
- 71. Interior Plateau\* (8.3.3)
  - 71m. Northern Shawnee Hills\*
  - 71n. Southern Shawnee Hills\*
- 72. Interior River Valleys and Hills (8.3.2)
  - 72a. Wabash / Ohio Bottomlands\*
  - 72d. Upper Mississippi Alluvial Plain
  - 72e. Middle Mississippi Alluvial Plain
  - 72f. River Hills
  - 72g. Southern Ozarkian River Bluffs\*
  - 72i. Western Dissected Illinoian Till Plain\*
  - 72j. Southern Illinoian Till Plain\*
  - 72k. Cretaceous Hills\*
  - 72l. Karstic Northern Ozarkian River Bluffs\*
- 73. Mississippi Alluvial Plain (8.5.2)
  - 73a. Northern Holocene Meander Belts
  - 73b. Northern Pleistocene Valley Trains\*
  - 73c. Saint Francis Lowlands
  - 73f. Western Lowlands Holocene Meander Belts
  - 73g. Western Lowlands Pleistocene Valley Trains
- 74. Valley Loess Plains (8.3.6)
  - 74a. Bluff Hills
  - 74b. Loess Plains

## Aquatic Subregions

Like terrestrial ecoregions, aquatic subregions can be classified in discrete units, based on watershed and hydrological characteristics. Subregions are composed of ecological drainage units, which are major watersheds that are consolidated based on similarity and proximity (MSDIS 2009a). Subregions and their component drainage units can be seen in Table 3 (below).

Naturally, many of the same patterns emerge in both terrestrial and aquatic geographic classifications since the two are not independent of one another. However, aquatic subregions and ecological drainage units can differ from their terrestrial counterparts since most aquatic organisms are confined to watersheds as well as specific habitats.

Since many species of amphibians and reptiles are partially or wholly dependent on aquatic habitats, distributions can sometimes be best understood and explained by examining aquatic subregions, ecological drainage units, and even individual watersheds.

The base map for major rivers and streams (page 12) shows locations of actual rivers and streams inside Missouri and just outside the state's borders where appropriate (USGS 1994; MDC 2007b). The watershed boundaries map (page 13) displays the maximum terrestrial boundary for major watersheds (NRCS 2002). Both of these maps are underlain with unique background colors to display Missouri's primary drainage systems: a) Mississippi River, b) Missouri River, c) Arkansas River, and d) White River.


Some species found in the state exhibit distributions that can be better understood by referring to these primary drainage systems. The ecological drainage units map (page 14) shows a somewhat consolidated view of the watersheds map based on those watersheds and other characteristics. It is underlain with unique background colors showing the three broad aquatic subregions in the state. Still more detailed aquatic region classifications are available but often include areas based on very small creeks and their watersheds and are therefore not included here.

**Table 3.** List of Aquatic Subregions and Ecological Drainage Units from MSDIS (2009a, 2009c). Each drainage unit is indicated with its primary drainage system: a) Mississippi River, b) Missouri River, c) Arkansas River, d) White River.

- ★ Central Plains Subregion
  - ★ Blackwater / Lamine (b)
  - ★ Cuivre / Salt (a)
  - ★ Des Moines (a)
  - ★ Grand / Chariton (b)
  - ★ Kansas (b)
  - ★ Nishnabotna / Platte (b)
  - ★ Osage / South Grand (b)
- ★ Ozarks Subregion
  - ★ Apple / Joachim (a)
  - ★ Black / Current (d)
  - ★ Gasconade (b)
  - ★ Meramec (a)
  - ★ Moreau / Loutre (b)
  - ★ Osage (b)
  - ★ Neosho (c)
  - ★ Upper Saint Francis / Castor (a)
  - ★ White (d)
- ★ Mississippi Alluvial Subregion
  - ★ Black / Cache (a)
  - ★ Saint Francis / Little (a)
  - ★ Saint Johns Bayou (a)

## Natural Divisions and Sections


Modified from Thom and Wilson (1980)  
Base map data provided by MSDIS (2009b) and MDC (2007a)




## Level I Terrestrial Ecoregions


Data provided by EPA (2003)

Base map data provided by the National Atlas (2001)










## Level IV Terrestrial Ecoregions


Data provided by EPA (2003)  
Base map data provided by the National Atlas (2001)



# Rivers and Major Streams


Data provided by USGS (1994) and MDC (2007b)  
Base map data provided by MSDIS (2009b)


- Mississippi River Watershed
- Missouri River Watershed
- White River Watershed
- Arkansas River Watershed






Data provided by NRCS (2002)  
Base map data provided by MSDIS (2009b)









## Species Locality Maps

Key:


- Museum or published record believed to be valid
- \* County record only for:
  - Valid record without a specific locality
  - Species of conservation concern, locality withheld
- ? Questionable record that may be valid




*Siren intermedia*  
Western Lesser Siren




*Cryptobranchus alleganiensis*  
Hellbender




*Ambystoma annulatum*  
Ringed Salamander




*Ambystoma maculatum*  
Spotted Salamander



*Ambystoma opacum*  
Marbled Salamander




*Ambystoma talpoideum*  
Mole Salamander




*Ambystoma texanum*  
Small-mouthed Salamander







*Ambystoma tigrinum*  
Eastern Tiger Salamander




*Notophthalmus viridescens*  
Central Newt




*Necturus maculosus*  
Mudpuppy




*Amphiuma tridactylum*  
Three-toed Amphiuma




*Eurycea longicauda*  
Long-tailed Salamander




*Eurycea lucifuga*  
Cave Salamander




*Eurycea spelaea*  
Grotto Salamander




*Eurycea tynerensis*  
Oklahoma Salamander




*Hemidactylium scutatum*  
Four-toed Salamander




*Plethodon albagula*  
Western Slimy Salamander




*Plethodon angusticlavius*  
Ozark Zigzag Salamander




*Plethodon serratus*  
Southern Red-backed Salamander




*Scaphiopus holbrookii*  
Eastern Spadefoot




*Spea bombifrons*  
Plains Spadefoot




*Anaxyrus americanus*  
American Toad




*Anaxyrus cognatus*  
Great Plains Toad




*Anaxyrus fowleri*  
Fowler's Toad




*Anaxyrus woodhousii*  
Woodhouse's Toad




*Acris blanchardi*  
Blanchard's Cricket Frog




*Hyla chrysoscelis*  
Cope's Gray Treefrog




*Hyla cinerea*  
Green Treefrog




*Hyla versicolor*  
Gray Treefrog




*Pseudacris crucifer*  
Spring Peeper




*Pseudacris feriarum*  
Upland Chorus Frog




*Pseudacris fouquettei*  
Cajun Chorus Frog




*Pseudacris illinoensis*  
Illinois Chorus Frog




*Pseudacris maculata*  
Boreal Chorus Frog




*Lithobates areolatus*  
Northern Crawfish Frog




*Lithobates blairi*  
Plains Leopard Frog



*Lithobates catesbeianus*  
American Bullfrog



*Lithobates clamitans*  
Green Frog




*Lithobates palustris*  
Pickerel Frog







*Lithobates pipiens*  
Northern Leopard Frog




*Lithobates sphenoccephalus*  
Southern Leopard Frog




*Lithobates sylvaticus*  
Wood Frog




*Gastrophryne carolinensis*  
Eastern Narrow-mouthed Toad




*Gastrophryne olivacea*  
Western Narrow-mouthed Toad




*Crotaphytus collaris*  
Eastern Collared Lizard




*Phrynosoma cornutum*  
Texas Horned Lizard




*Sceloporus consobrinus*  
Prairie Lizard




*Aspidoscelis sexlineata*  
Six-lined Racerunner




*Plestiodon anthracinus*  
Southern Coal Skink




*Plestiodon fasciatus*  
Common Five-lined Skink




*Plestiodon laticeps*  
Broad-headed Skink




*Plestiodon obsoletus*  
Great Plains Skink




*Plestiodon septentrionalis*  
Prairie Skink



*Scincella lateralis*  
Little Brown Skink




*Ophisaurus attenuatus*  
Western Slender Glass Lizard




*Hemidactylus turcicus*  
Mediterranean Gecko




*Podarcis siculus*  
Italian Wall Lizard




*Cemophora coccinea*  
Northern Scarletsnake




*Coluber constrictor*  
North American Racer




*Coluber flagellum*  
Eastern Coachwhip




*Lampropeltis calligaster*  
Prairie Kingsnake




*Lampropeltis holbrooki*  
Speckled Kingsnake




*Lampropeltis nigra*  
Eastern Black Kingsnake




*Lampropeltis triangulum*  
Eastern Milksnake




*Opheodrys aestivus*  
Northern Rough Greensnake




*Opheodrys vernalis*  
Smooth Greensnake




*Pantherophis emoryi*  
Great Plains Ratsnake




*Pantherophis obsoletus*  
Western Ratsnake



*Pantherophis ramspotti*  
Western Foxsnake




*Pantherophis vulpinus*  
Eastern Foxsnake




*Pituophis catenifer*  
Bullsnake







*Sonora semiannulata*  
Variable Groundsnake




*Tantilla gracilis*  
Flat-headed Snake




*Carphophis vermis*  
Western Wormsnake




*Diadophis punctatus*  
Ring-necked Snake




*Farancia abacura*  
Western Mudsucker




*Heterodon gloydi*  
Dusty Hog-nosed Snake




*Heterodon nasicus*  
Plains Hog-nosed Snake




*Heterodon platirhinos*  
Eastern Hog-nosed Snake




*Clonophis kirtlandii*  
Kirtland's Snake




*Haldea striatula*  
Rough Earthsnake




*Nerodia cyclopion*  
Mississippi Green Watersnake




*Nerodia erythrogaster*  
Plain-bellied Watersnake




*Nerodia fasciata*  
Broad-banded Watersnake




*Nerodia rhombifer*  
Diamond-backed Watersnake




*Nerodia sipedon*  
Common Watersnake




*Regina grahamii*  
Graham's Crawfish Snake




*Storeria dekayi*  
DeKay's Brownsnake




*Storeria occipitomaculata*  
Northern Red-bellied Snake




*Thamnophis proximus*  
Orange-striped Ribbonsnake




*Thamnophis radix*  
Plains Gartersnake




*Thamnophis sirtalis*  
Common Gartersnake




*Tropidoclonion lineatum*  
Lined Snake




*Virginia valeriae*  
Western Smooth Earthsnake




*Agkistrodon contortrix*  
Copperhead




*Agkistrodon piscivorus*  
Western Cottonmouth




*Crotalus horridus*  
Timber Rattlesnake




*Sistrurus catenatus*  
Eastern Massasauga




*Sistrurus miliarius*  
Western Pygmy Rattlesnake




*Sistrurus tergeminus*  
Prairie Massasauga



*Chelydra serpentina*  
Snapping Turtle




*Macrochelys temminckii*  
Alligator Snapping Turtle




*Kinosternon flavescens*  
Yellow Mud Turtle







*Kinosternon subrubrum*  
Mississippi Mud Turtle




*Sternotherus odoratus*  
Eastern Musk Turtle



*Chrysemys dorsalis*  
Southern Painted Turtle




*Chrysemys picta*  
Western Painted Turtle




*Deirochelys reticularia*  
Western Chicken Turtle




*Emydoidea blandingii*  
Blanding's Turtle




*Graptemys geographica*  
Northern Map Turtle




*Graptemys ouachitensis*  
Ouachita Map Turtle




*Graptemys pseudogeographica*  
False Map Turtle




*Pseudemys concinna*  
River Cooter




*Terrapene carolina*  
Three-toed Box Turtle




*Terrapene ornata*  
Ornate Box Turtle



*Trachemys scripta*  
Red-eared Slider



*Apalone mutica*  
Midland Smooth Softshell




*Apalone spinifera*  
Eastern Spiny Softshell


## County Records Maps

Key:


- Valid vouchered county record




*Siren intermedia*  
Western Lesser Siren




*Cryptobranchus alleganiensis*  
Hellbender




*Ambystoma annulatum*  
Ringed Salamander




*Ambystoma maculatum*  
Spotted Salamander




*Ambystoma opacum*  
Marbled Salamander




*Ambystoma talpoideum*  
Mole Salamander




*Ambystoma texanum*  
Small-mouthed Salamander




*Ambystoma tigrinum*  
Eastern Tiger Salamander




*Notophthalmus viridescens*  
Central Newt




*Necturus maculosus*  
Mudpuppy




*Amphiuma tridactylum*  
Three-toed Amphiuma




*Eurycea longicauda*  
Long-tailed Salamander



*Eurycea lucifuga*  
Cave Salamander




*Eurycea spelaea*  
Grotto Salamander




*Eurycea tynerensis*  
Oklahoma Salamander







*Hemidactylium scutatum*  
Four-toed Salamander




*Plethodon albagula*  
Western Slimy Salamander




*Plethodon angusticlavius*  
Ozark Zigzag Salamander




*Plethodon serratus*  
Southern Red-backed Salamander




*Scaphiopus holbrookii*  
Eastern Spadefoot




*Spea bombifrons*  
Plains Spadefoot




*Anaxyrus americanus*  
American Toad




*Anaxyrus cognatus*  
Great Plains Toad




*Anaxyrus fowleri*  
Fowler's Toad




*Anaxyrus woodhousii*  
Woodhouse's Toad




*Acris blanchardi*  
Blanchard's Cricket Frog




*Hyla chrysoscelis*  
Cope's Gray Treefrog




*Hyla cinerea*  
Green Treefrog




*Hyla versicolor*  
Gray Treefrog




*Pseudacris crucifer*  
Spring Peeper




*Pseudacris feriarum*  
Upland Chorus Frog




*Pseudacris fouquettei*  
Cajun Chorus Frog




*Pseudacris illinoensis*  
Illinois Chorus Frog




*Pseudacris maculata*  
Boreal Chorus Frog




*Lithobates areolatus*  
Northern Crawfish Frog




*Lithobates blairi*  
Plains Leopard Frog




*Lithobates catesbeianus*  
American Bullfrog




*Lithobates clamitans*  
Green Frog




*Lithobates palustris*  
Pickerel Frog




*Lithobates pipiens*  
Northern Leopard Frog




*Lithobates sphenoccephalus*  
Southern Leopard Frog




*Lithobates sylvaticus*  
Wood Frog




*Gastrophryne carolinensis*  
Eastern Narrow-mouthed Toad




*Gastrophryne olivacea*  
Western Narrow-mouthed Toad



*Crotaphytus collaris*  
Eastern Collared Lizard




*Phrynosoma cornutum*  
Texas Horned Lizard




*Sceloporus consobrinus*  
Prairie Lizard







*Aspidoscelis sexlineata*  
Six-lined Racerunner




*Plestiodon anthracinus*  
Southern Coal Skink




*Plestiodon fasciatus*  
Common Five-lined Skink




*Plestiodon laticeps*  
Broad-headed Skink




*Plestiodon obsoletus*  
Great Plains Skink




*Plestiodon septentrionalis*  
Prairie Skink



*Scincella lateralis*  
Little Brown Skink




*Ophisaurus attenuatus*  
Western Slender Glass Lizard




*Hemidactylus turcicus*  
Mediterranean Gecko




*Podarcis siculus*  
Italian Wall Lizard




*Cemophora coccinea*  
Northern Scarletsnake




*Coluber constrictor*  
North American Racer




*Coluber flagellum*  
Eastern Coachwhip




*Lampropeltis calligaster*  
Prairie Kingsnake




*Lampropeltis holbrooki*  
Speckled Kingsnake




*Lampropeltis nigra*  
Eastern Black Kingsnake




*Lampropeltis triangulum*  
Eastern Milksnake




*Opheodrys aestivus*  
Northern Rough Greensnake




*Opheodrys vernalis*  
Smooth Greensnake




*Pantherophis emoryi*  
Great Plains Ratsnake




*Pantherophis obsoletus*  
Western Ratsnake




*Pantherophis ramspotti*  
Western Foxsnake




*Pantherophis vulpinus*  
Eastern Foxsnake




*Pituophis catenifer*  
Bullsnake




*Sonora semiannulata*  
Variable Groundsnake




*Tantilla gracilis*  
Flat-headed Snake




*Carphophis vermis*  
Western Wormsnake




*Diadophis punctatus*  
Ring-necked Snake




*Farancia abacura*  
Western Mudsucker



*Heterodon gloydi*  
Dusty Hog-nosed Snake




*Heterodon nasicus*  
Plains Hog-nosed Snake




*Heterodon platirhinos*  
Eastern Hog-nosed Snake







*Clonophis kirtlandii*  
Kirtland's Snake




*Haldea striatula*  
Rough Earthsnake




*Nerodia cyclopion*  
Mississippi Green Watersnake




*Nerodia erythrogaster*  
Plain-bellied Watersnake




*Nerodia fasciata*  
Broad-banded Watersnake




*Nerodia rhombifer*  
Diamond-backed Watersnake




*Nerodia sipedon*  
Common Watersnake




*Regina grahamii*  
Graham's Crawfish Snake




*Storeria dekayi*  
DeKay's Brownsnake




*Storeria occipitomaculata*  
Northern Red-bellied Snake




*Thamnophis proximus*  
Orange-striped Ribbonsnake




*Thamnophis radix*  
Plains Gartersnake




*Thamnophis sirtalis*  
Common Gartersnake




*Tropidoclonion lineatum*  
Lined Snake




*Virginia valeriae*  
Western Smooth Earthsnake




*Agkistrodon contortrix*  
Copperhead




*Agkistrodon piscivorus*  
Western Cottonmouth




*Crotalus horridus*  
Timber Rattlesnake




*Sistrurus catenatus*  
Eastern Massasauga




*Sistrurus miliarius*  
Western Pygmy Rattlesnake




*Sistrurus tergeminus*  
Prairie Massasauga




*Chelydra serpentina*  
Snapping Turtle




*Macrochelys temminckii*  
Alligator Snapping Turtle




*Kinosternon flavescens*  
Yellow Mud Turtle




*Kinosternon subrubrum*  
Mississippi Mud Turtle




*Sternotherus odoratus*  
Eastern Musk Turtle




*Chrysemys dorsalis*  
Southern Painted Turtle




*Chrysemys picta*  
Western Painted Turtle




*Deirochelys reticularia*  
Western Chicken Turtle



*Emydoidea blandingii*  
Blanding's Turtle




*Graptemys geographica*  
Northern Map Turtle




*Graptemys ouachitensis*  
Ouachita Map Turtle







*Graptemys pseudogeographica*  
False Map Turtle




*Pseudemys concinna*  
River Cooter




*Terrapene carolina*  
Three-toed Box Turtle




*Terrapene ornata*  
Ornate Box Turtle



*Trachemys scripta*  
Red-eared Slider



*Apalone mutica*  
Midland Smooth Softshell



*Apalone spinifera*  
Eastern Spiny Softshell

## Acknowledgements

Data for the base county maps for the locality and county distributions, the labeled county map, and the ecological drainage units and aquatic subregions were generously provided by the Missouri Spatial Data Information Service (MSDIS 2009a, 2009b, 2009c). Data for the major rivers were provided by the United States Geological Survey (USGS 1994) and supplemented with data from the Missouri Department of Conservation (MDC 2007). Data for Missouri natural divisions and sections (based on Thom and Wilson (1980)) were generously provided by the Missouri Department of Conservation (MDC 2007). Data for watershed boundaries were provided by the Natural Resources Conservation Service (NRCS 2002). Data for ecoregions were provided by the US Environmental Protection Agency (EPA 2003). Base map data for surrounding states were provided by the National Atlas of the United States (National Atlas 2001).

All technology-related aspects of the Missouri Herpetological Atlas Project are supported by a suite of open source software that is both *gratis* (free as in beer) and *libre* (free as in speech). Kubuntu Linux is used as the operating system to host and execute all processes related to the project. Atlas, checklist, and GIS data are stored in a PostgreSQL database, which has been spatially enabled using PostGIS. Project documentation and database scripts are written with PHP and displayed using an Apache web server. QGIS is used to produce all base maps, distribution, county records, and other maps. The programming language Python is used to create GIS plug-ins for map exports. The open source Python library, ReportLab, is used to create the final atlas publication in portable document format (PDF), which was recently declared an ISO standard.

Most importantly, a project of this magnitude cannot be accomplished without the contributions of many individuals. We would like to thank the following for specimen information, feedback, and technical assistance on the project:

Nathan Aaron, Scott Altnether, Tom Anderson, Jonathan Ashley, Ralph Axtell, D. Baker, Stephanie Bartlett, B. Barton, Neil Bass, Tom Beauvais, Bill Becker, Don Becker, M. Bedford, R. Berry, A. Beston, Beth Blaisdell, Tyler Blake, Aaron Bossert, Michelle Bowe, Mark Bower, Andrew Braun, Jeff Briggler, Danny Brown, Sara Bruce, Josh Bruegge, Kevin Brunke, Carol Bryant, Steve Buback, Jenny Bunch, Frank Burbrink, John Cadle, T. Carpenter, Gary Casper, Rob Caviness, Ellen Censky, C. Chambers, D. Clark, Clint Coffey, Shayla Coffey, Skyler Collins, Matt Coombs, Wayne Corse, Danny Counts, C. Crabtree, Mike Cravens, Joe Cridlebaugh, Wyatt Cridlebaugh, R. Dames, Charles Dardin, Jeff Dawson, B. Day, B. DeCavele, J. Decker, Drew Dittmer, Scott Doherty, R. Douglas, Joseph Douglass, D. Drake, Frank Drummond, Frank Durbian, Eric East, David Easterla, Amber Edwards, T. Enright, P. Eyheralde, E. Field, Amanda Fisher, Z. Fratto, G. Frazier, Mark Frazier, Dillon Freiburger, Paul Frese, Darrell Frost, Bonnie Gardner, C. Gemming, Carl Gerhardt, Matt Gifford, M. Gihring, Brian Gilbert, Bob Gillespie, T. Girard, Brad

Glorioso, Steve Gotte, Millie Graham, Brian Greene, Greg Gremaud, Adam Guerich, Brad Hadley, Adam Hamlin, D. Harrison, Kevin Hedgpeth, A.J. Hendershott, Robert Henderson, Mary Hennen, John Hess, Joshua Hill, Dan Hocking, Colby Hodges, Dana Hoisington, Ethan Hollender, Terry Holtsclaw, E. Hooper, Emily Horner, Ann Horsman, Brian Hubbs, Jamie Hubert, James Hunt, Seth Ince, Adam Ivey, Thomas Jay, Matt Jeppson, Tom Jessen, Tom Johnson, Mike Jones, Jim Lamer, John Karel, Tim Kavan, Kate Kelly, Bob Kipfer, Arnold Kluge, Bob Krager, K. Korthas, Mike Leahy, Kevin Lewis, S. Liebi, Day Ligon, Elicia Ligon, Kenton Lohraff, Alexandria London, Peter Lowther, Amy McClure, Roy McDiarmid, Elizabeth McGhee, B. McKeage, Paul McKenzie, Don McKnight, Jennifer Maag, Glenn Manning, Nicholas March, Dexter Mardis, Gina Martin, Frank Martinez, Rebecca Maxfield, Christine Mayer, Arianne Messerman, Dean Metter, C. Meyer, Brian Miller, Danielle Miller, J. Miller, N. Mitchell, Chad Montgomery, D. Moore, T. Moore, L. Moree, Denise Morgan-Irish, K. Morrow, Sarah Moss, Nathan Muenks, Tom Nagel, Kevin Narum, L. Neuman-Lee, Chris Newbold, Abby Nicholson, Krista Noel, Laurel Norton, Jim Oehmke, Tyler Olcott, Matt Ormsby, Seth Otto, Cheryl Palmer, Peter Paplanus, John Par, Caleb Pemberton, Bill Peterman, M. Peterson, Katie Petrie, Tim Pettinelli, Chris Phillips, Nick Pieroni, G. Pinson, G. Pitchford, P. Pitts, J. Pollock, Crystal Poole, Robert Powell, Tracey Prenger, B. Probst, Keith Pulles, Holly Quinn, Hugh Quinn, Corey Raimond, Alan Resetar, Joe Ridings, Daren Riedle, Rhonda Rimer, Tony Robyn, Bernard Ross, Douglas Rossman, David Rostollan, Chad Runco, B. Russell, Greg Schneider, Tim Schoppenhorst, E. Schott, Bruce Schuette, Jeremy Schumacher, C. Scott, K. Scott, J. Severson, Chris Shelton, Chris Shulse, Dustin Siegel, Carri Silman, John Simmons, Rachel Simmons, Carl Smith, Geoff Smith, Isaac Smith, Madeline Smith, S. Smith, Steven Snow, Leon Spring, Marianne Stanley, Clarissa Starbuck, Andy Starostka, Shane Staten, B. Stehl, Jeff Stewart, Greg Stoner, James Studdard, K. Sulkowski, Becky Swearingen, Travis Taggart, Jeremiah Terrell, Wendell Tharp, Ryan Thies, M. Thompson, A. Thorn, Bart Tibbs, Mark Tovar, Sean Tracy, Stan Trauth, Angelina Trombley, T. Trombone, Linda Trueb, J. Twardowski, Chad Wallis, Sammy Wentz, J. West, Steve Westin, Amber Wiewel, Brian Willey, Bethany Williams, Linda Williams, Myki Williamson, Matt Wishard, Rod Wittenberg, N. Woodland, Nathan Woolery, Reese Worthington, Nico Yoch, Elizabeth Yohe

## Literature Cited

- Anderson, P. 1942. Amphibians and reptiles of Jackson County, Missouri. *Bulletin of the Chicago Academy of Science* 6(11): 203-220.
- Anderson, P. 1945. New herpetological records for Missouri. *Bulletin of Chicago Academy of Sciences* 7(5): 271-275.
- Anderson, P. 1957. A second list of new herpetological records for Missouri. *Natural History Miscellanea* (161): 1-5.
- Anderson, P. 1965. *The Reptiles of Missouri*. University of Missouri Press, Columbia, Missouri. xxiii + 330p.

- Blanchard, F.N. 1924. A collection of amphibians and reptiles from southeastern Missouri and Illinois. *Papers of the Michigan Academy of Science, Art, and Letters* 4: 533-541.
- Bodie, J.R. 1998. Contemporary hydrology of a large-river floodplain: Multiple-scale effects on freshwater turtles. Unpublished MA thesis, University of Missouri-Columbia.
- Bonett, R.M. and P.T. Chippindale. 2004. Speciation, phylogeography and evolution of life history and morphology in plethodontid salamanders of the *Eurycea multiplicata* complex. *Molecular Ecology* 13: 1189-1203.
- Brandley, M.C., A. Schmitz and T.W. Reeder. 2005. Partitioned Bayesian analyses, partition choice, and the phylogenetic relationships of Scincid lizards. *Systematic Biology* 54(3): pp. 373-390.
- Briggler, J.T. and T.R. Johnson. 2021. *The Amphibians and Reptiles of Missouri* (3rd ed.). Missouri Department of Conservation, Jefferson City, Missouri.
- Briggler, J.T., R.L. Rimer and G. Deichsel. 2015. First record of the Northern Italian Wall Lizard (*Podarcis siculus campestris*) in Missouri. *IRCF Reptiles and Amphibians* 22(1): 43-45.
- Boyer, D.A., and A.A. Heinze. 1934. An annotated list of the amphibians and reptiles of Jefferson County, Missouri. *Transactions of the Saint Louis Academy of Science* 28(4): 185-200.
- Buflino, A.P. 2004. Geographic Distribution: *Hemidactylus turcicus turcicus*. *Herpetological Review* 35: 188.
- Burke, R.L. and G. Deichsel. 2008. Lacertid lizards introduced into North America: History and future, pp. 347-353. In: J.C. Mitchell, R.E.I. Brown and B. Bartholomew (eds.) *Urban Herpetology*. Herpetological Conservation vol. 3. SSAR, Salt Lake City, Utah.
- Burt, C.E. 1932. Records of amphibians from the eastern and central United States (1931). *American Midland Naturalist* 13: 75-85.
- Burt, C.E. 1933. A collection of amphibians and reptiles from southern Missouri. *American Midland Naturalist* 14: 170-173.
- Burt, C.E. 1935. Further records of the ecology and distribution of amphibians and reptiles in the middle west. *American Midland Naturalist* 16(3): 311-366.
- Burt, C.E. and W.L. Hoyle. 1935. Additional records of the reptiles of the central prairie region of the United States. *Transactions of the Kansas Academy of Science* 37: 193-216.
- Conant, R. 1960. The Queen Snake, *Natrix septemvittata*, in the Interior Highlands of Arkansas and Missouri, with comments upon similar disjunct distributions. *Proceedings of The Academy of Natural Sciences of Philadelphia* 112(2): 25-40.
- Conant, R., and J.T. Collins. 1998. *A Field Guide to Reptiles and Amphibians: Eastern and Central North America*. (3rd ed., expanded) Peterson Field Guides, Houghton Mifflin Company, New York. 616p.
- Conant, R., and C.J. Goin. 1948. A new subspecies of the soft-shelled turtle from the central United States, with comments on the application of the name *Amyda*. *Occasional Papers of the Museum of Zoology* (510): 1-19.
- Crother, B.I. (chair, Committee on Standard English and Scientific Names) 2017. Scientific and Standard English Names of Amphibians and Reptiles of North America, North of Mexico, with Comments Regarding Confidence in our Understanding, 8th ed. SSAR *Herpetological Circular* (43): 102p.
- Crother, B.I., M.E. White, J.M. Savage, M.E. Eckstut, M.R. Graham, and D.W. Gardner. 2011. Reevaluation of the Status of the Foxsnakes *Pantherophis gloydi* Conant and *P. vulpinus* Baird and Girard (Lepidosauria). *ISRN Zoology* 2011 (Article ID 436049):15.
- Daniel, R.E. 2007. New size records for three species of Missouri reptiles. *Missouri Herpetological Association Newsletter* 20: 16-17.
- Daniel, R.E., and B.S. Edmond. 2000. New and previously unreported distribution records of amphibians and reptiles in Missouri for 2000. *Missouri Herpetological Association Newsletter* 13: 14-19.
- Daniel, R.E., and B.S. Edmond. 2001. New and previously unreported distribution records of amphibians and reptiles in Missouri for 2001. *Missouri Herpetological Association Newsletter* 14: 7-12.
- Daniel, R.E., and B.S. Edmond. 2002a. *Atlas of Missouri Amphibians and Reptiles (for 2001)*. <<http://atlas.moherp.org/pubs/atlas01.pdf>>
- Daniel, R.E., and B.S. Edmond. 2002b. Revised County Distribution Maps of Amphibians and Reptiles of Missouri. *Missouri Herpetological Association Newsletter* 15: 16-38.
- Daniel, R.E., and B.S. Edmond. 2003. *Atlas of Missouri Amphibians and Reptiles (for 2002)*. <<http://atlas.moherp.org/pubs/atlas02.pdf>>
- Daniel, R.E., and B.S. Edmond. 2004. *Atlas of Missouri Amphibians and Reptiles (for 2003)*. <<http://atlas.moherp.org/pubs/atlas03.pdf>>
- Daniel, R.E., and B.S. Edmond. 2005. *Atlas of Missouri Amphibians and Reptiles (for 2004)*. <<http://atlas.moherp.org/pubs/atlas04.pdf>>
- Daniel, R.E., and B.S. Edmond. 2006. *Atlas of Missouri Amphibians and Reptiles for 2005*. <<http://atlas.moherp.org/pubs/atlas05.pdf>>
- Daniel, R.E., and B.S. Edmond. 2008. *Atlas of Missouri Amphibians and Reptiles for 2007*. <<http://atlas.moherp.org/pubs/atlas07.pdf>>
- Daniel, R.E., and B.S. Edmond. 2010. *Atlas of Missouri Amphibians and Reptiles for 2009*. <<http://atlas.moherp.org/pubs/atlas09.pdf>>
- Daniel, R.E., and B.S. Edmond. 2012. *Atlas of Missouri Amphibians and Reptiles for 2011*. <<http://atlas.moherp.org/pubs/atlas11.pdf>>
- Daniel, R.E., and B.S. Edmond. 2013. *Atlas of Missouri Amphibians and Reptiles for 2012*. <<http://atlas.moherp.org/pubs/atlas12.pdf>>
- Daniel, R.E., and B.S. Edmond. 2014. *Atlas of Missouri Amphibians and Reptiles for 2013*. <<http://atlas.moherp.org/pubs/atlas13.pdf>>
- Daniel, R.E., and B.S. Edmond. 2017. *Atlas of Missouri Amphibians and Reptiles for 2016*. <<http://atlas.moherp.org/pubs/atlas16.pdf>>

- Daniel, R.E., and B.S. Edmond. 2018. *Atlas of Missouri Amphibians and Reptiles for 2017*. <<http://atlas.moherp.org/pubs/atlas17.pdf>>
- Daniel, R.E., and B.S. Edmond. 2019. *Atlas of Missouri Amphibians and Reptiles for 2018*. <<http://atlas.moherp.org/pubs/atlas18.pdf>>
- Daniel, R.E., and B.S. Edmond. 2020. *Atlas of Missouri Amphibians and Reptiles for 2019*. <<http://atlas.moherp.org/pubs/atlas19.pdf>>
- Daniel, R.E., and B.S. Edmond. 2021. *Atlas of Missouri Amphibians and Reptiles for 2020*. <<http://atlas.moherp.org/pubs/atlas20.pdf>>
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2002. New and previously unreported distribution records of amphibians and reptiles in Missouri for 2002. *Missouri Herpetological Association Newsletter* 15: 9-15.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2003. New and previously unreported herpetological records from Missouri for 2003. *Missouri Herpetological Association Newsletter* 16: 11-15.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2004. New herpetological records from Missouri for 2004. *Missouri Herpetological Association Newsletter* 17: 9-12.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2005. New herpetological records from Missouri for 2005. *Missouri Herpetological Association Newsletter* 18: 8-11.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2006. New herpetological records from Missouri for 2006. *Missouri Herpetological Association Newsletter* 19: 9-12.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2007. New herpetological records from Missouri for 2007. *Missouri Herpetological Association Newsletter* 20: 10-13.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2008. New herpetological records from Missouri for 2008. *Missouri Herpetological Association Newsletter* 21: 9-14.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2009. New herpetological distribution records for Missouri in 2009. *Missouri Herpetological Association Newsletter* 22: 7-9.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2010. New herpetological distribution records for Missouri in 2010. *Missouri Herpetological Association Newsletter* 23: 6-8.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2011. New herpetological distribution records for Missouri in 2011. *Missouri Herpetological Association Newsletter* 24: 14-16.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2012. New herpetological distribution records for Missouri in 2012. *Missouri Herpetological Association Newsletter* 25: 11-15.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2013. New herpetological distribution records for Missouri in 2013. *Missouri Herpetological Association Newsletter* 26: 11-15.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2014. New herpetological distribution records for Missouri in 2014. *Missouri Herpetological Association Newsletter* 27: 15-18.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2015. New herpetological distribution records for Missouri in 2015. *Missouri Herpetological Association Newsletter* 28: 10-14.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2016. New herpetological distribution records for Missouri in 2016. *Missouri Herpetological Association Newsletter* 29: 16-19.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2017. New herpetological distribution records for Missouri in 2017. *Missouri Herpetological Association Newsletter* 30: 10-12.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2018. New herpetological distribution records for Missouri in 2018. *Missouri Herpetological Association Newsletter* 31: 11-13.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2019. New herpetological distribution records for Missouri in 2019. *Missouri Herpetological Association Newsletter* 32: 8-10.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2020. New herpetological distribution records for Missouri in 2020. *Missouri Herpetological Association Newsletter* 33: 3-11.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2021. New herpetological distribution records for Missouri in 2021. *Missouri Herpetological Association Newsletter* 34: 3-6.
- Daniel, R.E., B.S. Edmond, and T.R. Johnson. 1998. New and previously unreported records of amphibians and reptiles in Missouri for 1998. *Missouri Herpetological Association Newsletter* 11: 8-17.
- Daniel, R.E., B.S. Edmond, and T.R. Johnson. 1999. New and previously unreported records of amphibians and reptiles in Missouri for 1999. *Missouri Herpetological Association Newsletter* 12: 10-18.
- Easterla, D.A. 1972. Herpetological records for northwest Missouri. *Transactions of the Missouri Academy of Science* 6: 158-160.
- Easterla, D.A. and M.D. Meadows. 1993. The northern prairie skink rediscovered in Missouri, the last recorded occurrence of the smooth green snake for Missouri, plus a Missouri range extension for the northern spring peeper. *Bluebird* 60(3): 105-109.
- Edmond, B.S. and R.E. Daniel. 2022. *Missouri Herpetological Atlas Project*. <<http://atlas.moherp.org/>> Accessed 03 August 2022.
- Edmond, B.S. and R.E. Daniel. 2014. A New Kingsnake for Missouri and Some Comments on the Biogeography of Southeast Missouri. *Missouri Herpetological Association Newsletter* 27: 18-21.
- Environmental Protection Agency (EPA). 2003. *Level III and IV Ecoregions of the Continental United States*. <[http://www.epa.gov/wed/pages/ecoregions/na\\_eco.htm](http://www.epa.gov/wed/pages/ecoregions/na_eco.htm)> Accessed 1 December 2009.
- Evans, P.D. 1940. Notes on Missouri snakes. *Copeia* 1940: 53-54.

- Frost, D., T. Grant, J. Faivovich, R. Bain, A. Haas, C. Haddad, R. De Sá, A. Channing, M. Wilkinson, S. Donnellan, C. Raxworthy, J. Campbell, B. Blotto, P. Moler, R. C. Drewes, R. Nussbaum, J. Lynch, D. Green and W. Wheeler. 2006. The amphibian tree of life. *Bulletin of the American Museum of Natural History* (297): pp. 1-370.
- Gamble, T., P.B. Berendzen, H.B. Shaffer, D.E. Starkey, and A.M. Simons. 2008. Species limits and phylogeography of North American Cricket Frogs (*Acris*: Hylidae). *Molecular Phylogenetics and Evolution* 48: 112-125.
- Gibbs, H.L., Murphy, M., and J.E. Chiucchi. 2011. Genetic identity of endangered massasauga rattlesnakes (*Sistrurus* sp.) in Missouri. *Conservation Genetics* 12: 433-439.
- Hawker, J.L. 1992. *Missouri Landscapes: A Tour Through Time*. Missouri Department of Natural Resources, Division of Geology and Land Survey, Rolla. 326pp.
- Henning, W.L. 1938. Amphibians and reptiles of a 2,220-acre tract in central Missouri. *Copeia* 1938(2): 91-92.
- Hubbs, B. 2011. Geographic Distribution: *Lampropeltis triangulum sypila*. *Herpetological Review* 42: 114.
- Hubbs, B. 2012. Geographic Distribution: *Apalone mutica*. *Herpetological Review* 43(3): 441.
- Hurter, J. 1911. Herpetology of Missouri. *Transactions of the Saint Louis Academy of Science* 20(5): 59-274.
- Johnson, T.R. 1987. *The Amphibians and Reptiles of Missouri*. (1st ed.) Missouri Department of Conservation, Jefferson City, Missouri. 368pp.
- Johnson, T.R. 2000. *The Amphibians and Reptiles of Missouri*. (2nd ed.) Missouri Department of Conservation, Jefferson City, Missouri. 400p.
- Johnson, T.R. and R.N. Bader. 1974. Annotated checklist of Missouri amphibians and reptiles. *St. Louis Herpetological Society Special Issue* 1: 1-16.
- Johnson, T.R. and R. Powell. 1988. New records of amphibians and reptiles in Missouri for 1988. *Missouri Herpetological Association Newsletter* 1: 4-8.
- Jones, J.M. 1967. A western extension of the known range of Kirtland's watersnake. *Herpetologica* 23(1): 66-67.
- Karel, J.A. 1974. Cottonmouths in the St. Francois Mountains. *Saint Louis Herpetological Society Newsletter* 1(8): 10.
- Kubatko, Laura S., Gibbs, H. Lisle, and Erik W. Bloomquist. 2011. Inferring Species-Level Phylogenies and Taxonomic Distinctiveness Using Multilocus Data in *Sistrurus* Rattlesnakes. *Systematic Biology* 60(4): 393-409.
- Leaché, A.D. and T.W. Reeder. 2002. Molecular systematics of the Eastern Fence Lizard (*Sceloporus undulatus*): A comparison of parsimony, likelihood, and Bayesian approaches. *Systematic Biology* 51(1): 44-68.
- Lemmon, E.M., A.R. Lemmon, J.T. Collins, J.A. Lee-Yaw and D.C. Cannatella. 2007. Phylogeny-based delimitation of species boundaries and contact zones in the trilling chorus frogs (*Pseudacris*). *Molecular Phylogenetics and Evolution* 44: 1068-1082.
- Lemmon, E.M., A.R. Lemmon, J.T. Collins and D.C. Cannatella. 2008. A New North American Chorus Frog Species (*Pseudacris*: Hylidae: Amphibia) from the South-Central United States. *Zootaxa*. 1675: 1-30.
- McConkey, E.H. 1954. A systematic study of the North American lizards of the genus *Ophisaurus*. *American Midland Naturalist* 51(1): 133-171.
- Missouri Department of Conservation (MDC). 2007a. *Natural Divisions and Sections*. <<http://www.mdc.mo.gov/>> Accessed 1 February 2007.
- Missouri Department of Conservation (MDC). 2007b. *Rivers and Streams*. <<http://www.mdc.mo.gov/>> Accessed 1 February 2007.
- Missouri Spatial Data Information Service (MSDIS). 2009a. *Aquatic Subregions*. <<http://msdis.missouri.edu/>> Accessed 1 December 2009.
- Missouri Spatial Data Information Service (MSDIS). 2009b. *Base County and Boundary Map*. <<http://msdis.missouri.edu/>> Accessed 1 December 2009.
- Missouri Spatial Data Information Service (MSDIS). 2009c. *Ecological Drainage Units*. <<http://msdis.missouri.edu/>> Accessed 1 December 2009.
- Myers, C.W. 1957. Amphibians and reptiles of Washington State Park, Washington County, Missouri. *Transactions of the Kansas Academy of Science* 60(3): 288-293.
- Myers, C.W. 1959. Amphibians and reptiles of Montauk State Park and vicinity, Dent County, Missouri. *Transactions of the Kansas Academy of Science* 62(1): 88-90.
- Nagy, Z. T., R. Lawson, U. Joger and M. Wink . 2004. Molecular systematics of racers, whipsnakes and relatives (Reptilia: Colubridae) using mitochondrial and nuclear markers. *Journal of Zoological Systematics and Evolutionary Research* 42: pp. 223-233.
- National Atlas of the United States. 2001. *State and County Boundaries of the United States*. <<http://nationalatlas.gov/atlasftp.html>> Accessed 1 December 2009.
- Natural Resources Conservation Service, U.S. Department of Agriculture. 2002. *Watershed Boundaries of the United States*. <<http://datagateway.nrcs.usda.gov/>> Accessed 1 December 2009.
- Nelson, P.W. 1987. *The Terrestrial Natural Communities of Missouri*. (revised ed.) Missouri Natural Areas Committee, Jefferson City. 197pp.
- Nelson, P.W. 2005. *The Terrestrial Natural Communities of Missouri*. Missouri Natural Areas Committee, Jefferson City. 550pp.
- Nicholson, A.E. 2013a. Geographic distribution: USA: Missouri: *Notophthalmus viridescens*. *Herpetological Review* 44(3): 470.
- Nicholson, A.E. 2013b. Geographic distribution: USA: Missouri: *Plestiodon laticeps*. *Herpetological Review* 44(3): 475.
- Nicholson, A.E. 2013c. Geographic distribution: USA: Missouri: *Lampropeltis triangulum*. *Herpetological Review* 44(3): 477.
- Nickerson, M.A., and R. Krager. 1971. Noteworthy records of Missouri reptiles. *Transactions of the Kansas Academy of Science* 74(1): 99-101.

- Nickerson, M.A. and R. Krager. 1972. Additional noteworthy records of Missouri amphibians and reptiles with a possible addition to the herpetofauna. *Transactions of the Kansas Academy of Science* 75(3): 276-277.
- Nigh, T.A., and W.A. Schroeder. 2002. *Atlas of Missouri Ecoregions*. Missouri Department of Conservation, Jefferson City. 212pp.
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. Map Supplement (scale 1:7,500,000). *Annals of the Association of American Geographers* 77(1): 118-125.
- Owens, V. 1949. New snake records and notes, Morgan County, Missouri. *Herpetologica* 5: 49-50.
- Peters, J.A. 1946. Reptiles and amphibians of Sam A. Baker State Park, Wayne County, Missouri. *Copeia* 1946(1): 44.
- Peters, J.A. 1946. Records of certain North American salamanders. *Copeia* 1946(2): 106.
- Pflieger, W.L. 1989. Aquatic community classification system for Missouri. *Aquatic Series No. 19*. Missouri Department of Conservation, Jefferson City.
- Phillips, C.A. 1994. Geographic distribution of mitochondrial DNA variants and the historical biogeography of the spotted salamander, *Ambystoma maculatum*. *Evolution* 48(3): 597-607.
- Powell, R., R. Conant, and J.T. Collins. 2016. *Peterson Field Guide to Reptiles and Amphibians of Eastern and Central North America*. (4th ed.) Houghton Mifflin Harcourt Publishing Company, New York. 494p.
- Powell, R., T.R. Johnson, and J.S. Parmelee. 1993. Updated distribution maps for amphibians and reptiles in Missouri. *Missouri Herpetological Association Newsletter* 6: 7-26.
- Powell, R., T.R. Johnson, and D.D. Smith. 1989. New records of amphibians and reptiles in Missouri for 1989. *Missouri Herpetological Association Newsletter* 2: 4-8.
- Powell, R., T.R. Johnson, and D.D. Smith. 1990. New records of amphibians and reptiles in Missouri for 1990. *Missouri Herpetological Association Newsletter* 3: 4-8.
- Powell, R., T.R. Johnson, and D.D. Smith. 1991. New records of amphibians and reptiles in Missouri for 1991. *Missouri Herpetological Association Newsletter* 4: 5-10.
- Powell, R., T.R. Johnson, and D.D. Smith. 1992. New records of amphibians and reptiles in Missouri for 1992. *Missouri Herpetological Association Newsletter* 5: 6-13.
- Powell, R., T.R. Johnson, and D.D. Smith. 1993. New records of amphibians and reptiles in Missouri for 1993. *Missouri Herpetological Association Newsletter* 6: 3-7.
- Powell, R., T.R. Johnson, and D.D. Smith. 1994. New records of amphibians and reptiles in Missouri for 1994. *Missouri Herpetological Association Newsletter* 7: 5-7.
- Powell, R., T.R. Johnson, and D.D. Smith. 1995. New records of amphibians and reptiles in Missouri for 1995. *Missouri Herpetological Association Newsletter* 8: 9-12.
- Powell, R., T.R. Johnson, and D.D. Smith. 1996. New records of amphibians and reptiles in Missouri for 1996. *Missouri Herpetological Association Newsletter* 9: 9-14.
- Powell, R., T.R. Johnson, and D.D. Smith. 1997. New records of amphibians and reptiles in Missouri for 1997. *Missouri Herpetological Association Newsletter* 10: 8-12.
- Powell, R. and R. Daniel. 1997. County distribution records of amphibians and reptiles in Missouri. *Missouri Herpetological Association Newsletter* 10: 12-69.
- Pyrone, R.A., and F.T. Burbrink. 2009. Systematics of the Common Kingsnake (*Lampropeltis getula*; Serpentes: Colubridae) and the burden of heritage in taxonomy. *Zootaxa* 2241: 22-32.
- Reeder, T.W., C.J. Cole, and H.C. Dessauer. 2002. Phylogenetic relationships of Whiptail lizards of the genus *Cnemidophorus* (Squamata: Teiidae): A test of monophyly, reevaluation of karyotypic evolution, and review of hybrid origins. *American Museum Novitates* (3365): 1-61.
- Shulse, C. 2006. Rediscovery of Kirtland's Snake (*Clonophis kirtlandii*) in Missouri. *Missouri Herpetological Association Newsletter* 19: 13-14.
- Shulse, C. 2007. Geographic Distribution: *Clonophis kirtlandii*. *Herpetological Review* 38: 220.
- Shaffer, H.B., and M.L. McKnight. 1996. The polytypic species revisited: Genetic differentiation and molecular phylogenetics of the tiger salamander *Ambystoma tigrinum* (Amphibia: Caudata) complex. *Evolution* 50(1): 417-433.
- Shepard, D.B. and A.R. Kuhns. 2017. Shifting rivers and stationary ground: biogeographic history of slimy salamanders on an isolated bluff in the Mississippi River floodplain. *Journal of Zoology* DOI:10.1111/jzo.12514
- Starkey, D.E., R.L. Burke, M.R.J. Forstner, J.B. Iverson, F.J. Janzen, A.G.J. Rhodin, G.R. Ultsch and H.B. Shaffer. 2003. Molecular systematics and phylogeography of the painted turtle (*Chrysemys picta*) complex. *Evolution* 57: 119-128.
- Steyermark, J.A. 1963. *Flora of Missouri*. Iowa State University Press, Ames. 1728pp.
- Tanner, W.W., and R.B. Loomis. 1957. Taxonomic and distributional study of the western subspecies of the milksnake, *Lampropeltis dolia*. *Transactions of the Kansas Academy of Science* 60(1): 12-42.
- Thom, R.H., and J.H. Wilson. 1980. The natural divisions of Missouri. *Transactions of the Missouri Academy of Science* 14:9-23.
- Thornhill, S.G. 1990. Genetic variation among populations of *Eurycea tynerensis* and neotenic *Eurycea multiplicata griseogaster*. Unpublished MS Thesis. Southwest Missouri State University, Springfield, Missouri.
- United States Geological Survey. 1994. *Major Rivers and Streams of the United States*. <<http://pubs.usgs.gov/>> Accessed 1 December 2009.
- Unklesbay, A.G., and J.D. Vineyard. 1992. *Missouri Geology: Three Billion Years of Volcanoes, Seas, Sediments, and Erosion*. University of Missouri Press, Columbia. 189pp.
- Utiger, U., N. Helfenberger, B. Schatti, C. Schmidt, M. Rufand, and V. Ziswiler. 2002. Molecular systematics

- and phylogeny of Old and New World rat snakes, *Elaphe* Auct., and related genera (Reptilia, Squamata, Colubridae). *Russian Journal of Herpetology* 9(2): 105-124.
- Van Wagenen, G. 1917. *Diadophis amabilis* in Missouri. *Copeia* (43): 43-44.
- Watkins, L.C. 1969. A third record of the four-toed salamander, *Hemidactylium scutatum*, in Missouri. *Transactions of the Kansas Academy of Science* 72(2): 264-265.
- Watkins, L.C., and L.L.Hinesley. 1970. Notes on the distribution and abundance of the sonoran skink, *Eumeces obsoletus*, in western Missouri. *Transactions of the Kansas Academy of Science* 73(1): 118-119.
- Williams, K.L. 1978. American Milk Snake. *Milwaukee Public Museum Publications in Biology and Geology* 2: 1-258.
- Yatskievych, G. 1999. *Steiermark's Flora of Missouri, Volume 1*. Missouri Department of Conservation, Jefferson City. 991pp.



## Appendix A: Sources

American Museum of Natural History  
Arkansas State University  
Austin Peay State University  
Bobby Witcher Memorial Collection, Avila College (KU)  
Carnegie Museum of Natural History  
Chicago Academy of Sciences  
Cornell University Museum of Vertebrates  
Culver-Stockton College  
Dallas Museum of Natural History  
Field Museum of Natural History  
H. Carl Gerhardt Audio Recording Collection (Cornell)  
Historical Literature Sources  
Illinois Natural History Survey  
Julius Hurter (Trans. St. Louis Acad. Sci 20(5): 59-274)  
Milwaukee Public Museum  
Missouri Department of Conservation  
Missouri State University  
Museum of Comparative Zoology, Harvard University  
Museum of Natural Science, Louisiana State University  
Museum of Southwest Biology (New Mexico)  
National Museum of Natural History  
Natural History Museum of Los Angeles County  
Paul Anderson (The Reptiles of Missouri)  
Pittsburg State University  
Saint Louis Academy of Sciences  
Southeast Missouri State University  
Southern Illinois University--Carbondale  
Southern Illinois University--Edwardsville  
Sternberg Museum of Natural History, Fort Hays State University  
Texas Natural History Collection  
University of Arkansas--Fayetteville  
University of California--Berkeley Museum of Vertebrate Zoology  
University of Central Missouri  
University of Florida at Gainesville  
University of Georgia, Museum of Natural History  
University of Illinois Natural History Museum  
University of Kansas, Museum of Natural History  
University of Michigan, Museum of Zoology  
University of Missouri--Columbia Dean E. Metter Memorial Herpetology Collection  
William Jewell College

\* Sources marked with an asterisk (\*) are included from published accounts only and are presumed to represent only part of the collection's Missouri records. All Missouri material from these sources has not been included in the database or the printed atlas.