# Atlas of Missouri Amphibians and Reptiles for 2023



Richard E. Daniel and Brian S. Edmond

# Atlas of Missouri Amphibians and Reptiles for 2023

Richard E. Daniel
Division of Biological Sciences
University of Missouri-Columbia
114 LeFevre Hall
Columbia MO 65211
danielr@missouri.edu

Brian S. Edmond Computer Services Missouri State University 901 South National Ave Springfield MO 65897 BrianEdmond@missouristate.edu

### Recommended citation:

Daniel, R.E. and B.S. Edmond. 2024. Atlas of Missouri Amphibians and Reptiles for 2023. <a href="http://atlas.moherp.org/pubs/atlas23.pdf">http://atlas.moherp.org/pubs/atlas23.pdf</a>

Updated: 19 April 2024 12:49

Copyright (C) 1997-2024. All rights reserved.

#### Table of Contents

## **Table of Contents**

Introduction
Amphibians and Reptiles of Possible Occurrence
Erroneously Reported and Non-Native Species
Distributions
Terrestrial Ecoregions
Aquatic Subregions
Natural Divisions and Sections Map
Level I Terrestrial Ecoregions Map
Level II Terrestrial Ecoregions Map
Level III Terrestrial Ecoregions Map
Level IV Terrestrial Ecoregions Map
Rivers and Major Streams Map
Watershed Boundaries Map
Ecological Drainage Units Map
County and Surrounding States Map
Species Locality Maps
County Records Maps
Acknowledgements
Literature Cited
Appendix A: Collection Sources
Appendix B: Literature Sources

### Introduction

Since 1988, members of the Missouri Herpetological Association have compiled new county distribution records for amphibians and reptiles native to the state (Daniel et al. 2022 and previous updates). Cumulative accounts of these new records were presented in Powell et al.. (1993), Powell and Daniel (1997), and Daniel and Edmond (2002). Johnson (2000) presented revised distribution maps based on a combination of these cumulative accounts and more recent annual compilations. The publication of Johnson (1987, 2000) renewed interest in expanding our knowledge of the distribution of the state's herpetofauna. Briggler and Johnson (2021) published an updated version of the The Amphibians and Reptiles of Missouri that incorporated all Atlas records up until that time. We follow their taxonomy and nomenclature here. In order to provide herpetologists working in Missouri with current information that is easily taken into the field, we are presenting revised species distribution and county records maps in a format that can be easily updated as new records are documented.

In 1997, with the initiation of the Missouri Herpetological Atlas Project (MOHAP), a database was established to serve as the basis for verifying new distribution records and tracking changes in individual species distributions within the state (Edmond and Daniel 2024). Qualification for inclusion in the database requires a catalogued voucher specimen housed in an institutional collection (Appendix A). However, in order to be valuable in tracking changes in the distribution of individual species, it is important to document historical records. Many of the specimens collected prior to the 1970s, most notably those reported by Hurter (1911) and Anderson (1965), were documented by specimens that have since been lost or destroyed. In order to provide the most accurate picture of the distribution of the state's herpetofauna we have included some literature records (Appendix B).

Some records were not included because the identification could not be determined with certainty. Within Missouri, the *Hyla versicolor* complex consists of two broadly sympatric and morphologically indistinguishable species (*H. chrysoscelis* and *H. versicolor*). Specimens collected for which the species identification was not determined by some non-morphological trait were not included.

Currently, the MOHAP database contains 38,487 entries and 34,349 valid, non-duplicated collections. This represents specimens housed in 37 museum collections and cited from 33 historical literature sources; 5,347 documented county records; 10,816 unique localities; and 18,150 unique species / locality combinations. This new total constitutes 679 new database entries since the 2021 edition of the *Atlas* (Daniel and Edmond 2022). The total number of species contained in the current edition of the atlas is 118, comprised of 116 native species and 2 non-native species.

Previous editions of the atlas are available upon request. Readers may contribute new records and read more about project details online (Edmond and Daniel 2024). Questions, comments, and suggestions should be directed to the senior author.

## Amphibians and Reptiles of Possible Occurrence

A number of species found in surrounding states are known to occur in close proximity to the borders of Missouri. While not currently recognized as part of the Missouri herpetofauna, some or all of these species may eventually be found within the state.

A number of primarily eastern species have distributions that reach the Mississippi River along a portion of the Missouri border. These include: Southern Two-lined Salamander (Eurycea cirrigera), Three-lined Salamander (Eurycea guttolineata), Northern Zigzag Salamander (Plethodon dorsalis), Northern Slimy Salamander (Plethodon glutinosus), Eastern Cricket Frog (Acris crepitans), Western Chorus Frog (Pseudacris triseriata), Bird-voiced Treefrog (Hyla avivoca), Eastern Fence Lizard (Sceloporus undulatus), Gray Ratsnake (Pantherophis spiloides), and Eastern Ribbonsnake (Thamnophis sauritus).

Rivers, even large ones, change course over time and typically form an imperfect geographic barrier. Shepard and Kuhns (2017) examined the separation between Northern Slimy Salamanders (*Plethodon glutinosus*) and Western Slimy Salamanders (*Plethodon albagula*) along the Illinois-Missouri border. They found that salamanders on Fountain Bluff, an island formed of Missouri uplands cut off by post-glacial changes in the Mississippi River channel, belong to the species found in Illinois.

Spotted Dusky Salamander (*Desmognathus conanti*) and Midwestern Earthsnake (*Carphophis amoenus*) have been reported in the southern portion of Crowley's Ridge of Arkansas. Crowley's Ridge is a thin, sandy upland formed during the Pleistocene, which extends from northeastern Arkansas through southeast Missouri to the Shawnee Hills of Illinois. Populations of one or both of these species may exist in the isolated remnants of Crowley's Ridge in Missouri.

Powell *et al.* (2016) shows the distribution of the Western Milksnake (*Lampropeltis gentilis*) extending into western Missouri along the Kansas border. However, the sparsity of records from this region makes it difficult to accurately define the species boundary. Until additional material is available for evaluation, we have not included this species as part of the state herpetofauna.

# Erroneously Reported and Non-Native Species

Anderson (1945) reported two specimens of the Dwarf Salamander (*Eurycea quadridigitata*) from Roaring River State Park in Barry County. With the nearest known naturally occurring populations of *E. quadridigitata* in southern Arkansas, more than 125 miles from the Missouri border, it is likely that this record is based on misidentified, but morphologically similar Oklahoma Salamanders (*E. tynerensis*).

Johnson and Bader (1974) included Lesser Earless Lizard (*Holbrookia maculata*) based on two specimens

collected from Knob Noster State Park in Johnson County. Nickerson and Krager (1972) considered this record problematic because of the close proximity of the park to Whiteman Air Force Base and Central Missouri State University. The nearest record of this species is more than 115 miles away in central Kansas. Further searches failed to produce additional specimens. As a result, this species is not considered to be part of Missouri's herpetofauna.

Anderson (1957) listed the occurrence of the Queensnake (*Regina septemvittata*) in Missouri based on three specimens deposited in the American Museum of Natural History collection. These specimens were ascribed to G.K. Noble and reported to come from Stone County. In further investigation by Roger Conant (1960), Byron C. Marshall, who worked with Noble in the Interior Highlands during this period, reportedly had no recollection of finding this species. The nearest known populations of Queen Snake are in the Boston Mountains of Arkansas, approximately 80 miles south of this locality. The questions about the origin of the specimens and the absence of additional records from Missouri suggest that this species does not occur in the state.

One species of non-native frog and two species of lizards have been reported in Missouri. A population of Greenhouse Frogs (*Eleutherodactylus planirostris*) was established inside a commercial greenhouse in Jefferson City, Missouri (J. Briggler, pers. comm.). The population of these small terrestrial breeding frogs persisted for several years, but has apparently disappeared following renovation of the greenhouse. There are no other reports of this tropical species occurring within the state. Its apparent inability to survive extreme winter weather makes it unlikely for these frogs to exist in Missouri as a free-ranging species outside of sheltered habitats, such as greenhouses.

Mediterranean Gecko (*Hemidactylus turcicus*) is native to the Mediterranean basin and western Asia. This species is easily transported and a highly successful colonizer. Populations of this species have been established worldwide. In the United States, this species is currently established widely throughout the south, especially along the Gulf and East Coasts. Isolated populations are also found around numerous urban centers of the Midwest and central plains (Powell *et al.* 2016). Two populations appear to be established in Missouri. Bufalino (2004) reported this species from St. Louis County. A second population was discovered in Joplin, Missouri by Mr. A. Braun (Daniel *et al.* 2015).

Briggler *et al.* (2015) reported the presence of a robust population of Italian Wall Lizard (*Podarcis siculus*) in a Joplin neighborhood after being contacted by a resident in 2013. This population apparently descended from a series of lizards imported from Topeka, KS, which escaped in 2001. In many states, including Missouri, it is illegal to release non-native species. Despite prohibitions, the majority of introduced populations of this species in North America originated from released animals in the pet trade (Burke and Deichsel 2008).

Thus far, non-native amphibians and reptiles in Missouri have been confined to urban centers and do not appear to constitute a threat to our native herpetofauna. However, this is not the case everywhere and non-native

species that have become invasive are considered by many biologists to be a major threat, second only to habitat loss or degradation, to native species. Responsible pet owners should always exercise care to prevent the spread of this and all non-native species into the natural environment.

### Distributions

Distribution is one of many characteristics used to describe a species and range maps (such as those found in Conant and Collins (1998)) are used to approximate a species' likely distribution. This atlas uses locality dot maps, with each dot representing a known locality for that species. This conservative method results in an underestimate of a species distribution but is more accurate than a range map. The purpose of the atlas project is to document as closely as possible both current and historical distributions for all native amphibians and reptiles in Missouri.

However, the printed version of the atlas only reflects the best historical approximation of each species' distribution in the state. The most noticeable resulting incongruity is that fact that some species are shown in historical localities in which they are almost certainly no longer found. For example, the Smooth Greensnake (Opheodrys vernalis) was once found in scattered populations in the prairie regions of the state. Due to extensive habitat loss and possibly other reasons, they are extremely rare or extirpated in the state.

While natural communities have long been a staple in ecology, formal classification of natural communities has been undertaken relatively recently in Missouri (Nelson 2005). Like individual species, natural communities can be described and characterized with distinct distributions and abundances in different ecoregions (see next section). For example, glades are common in the Ozarks but rare elsewhere in the state. Steyermark (1963), Yatskievych (1999), Pflieger (1989), Hawker (1992), and Unklesbay and Vineyard (1992) all describe and summarize the complex interplay among geology, natural history, ecoregions, natural divisions, and natural communities from slightly different perspectives. Taken together, these ideas form the basis of the biogeography of Missouri and they are essential in understanding why plants and animals have the distributions that we see today.

### Terrestrial Ecoregions

An ecoregion classification system attempts to define and describe geographic regions that correspond to broad ecosystem patterns, topography, geology, soils, vegetation patterns, and the distributions of plants and animals. Omernik (1987) described ecoregions (Levels I, II, and III) for the conterminous United States as a hierarchical scheme with Level I corresponding to large regions and Level III representing smaller, more precisely described regions. The Environmental Protection Agency (2003) is coordinating an effort to further subdivide Level III regions into Level IV regions. Nigh and Schroeder (2002) published Level III and Level IV ecoregions for Missouri.

Some amphibian and reptile species follow defined ecoregions closely. For example, the Cave Salamander (*Eurycea lucifuga*) is neatly confined to the Ozark Highlands (Level III) and found throughout the ecoregion, with the exception of most of the Springfield Plateau (Level IV). Several species found in the southeastern alluvial plain are particularly characteristic and also confined to that area (*e.g.*, Three-toed Amphiuma (*Amphiuma tridactylum*) and Southern Watersnake (*Nerodia fasciata*)).

Many species, however, seem to be abundant throughout Missouri, regardless of the region or natural community. The American Bullfrog (Lithobates catesbeianus), Snapping Turtle (Chelydra serpentina), and Western Ratsnake (Pantherophis obsoletus) among several others fall into this category. Finally, many species are more closely associated with a particular natural community or habitat than with a particular ecoregion or natural division. For example, The Flat-headed Snake (Tantilla gracilis) can be found in the Ozark Highlands, Osage Prairie, and Interior River Valleys and Hills but it is restricted to rocky glades.

In general, Level IV ecoregions are too granular to be useful in describing the distributions of Missouri amphibians and reptiles, several Ozark salamander species and many of the coastal plains species being notable exceptions. However, this level of detail is useful to understand natural community distributions on which some species are dependent. It also underscores the importance of looking at border states and across artificial political boundaries to understand distributions within Missouri.

Level I (page 8), Level II (page 9), Level III (page 10), and Level IV (page 11) ecoregion maps for Missouri and surrounding states, based on Omernik (1987) and Nigh and Schroeder (2002) are included here. The hierarchy for Levels I, II, and III is included in Table 1 (below), while the Level III and Level IV hierarchy is included in Table 2 (below). Unfortunately, the authors' coding scheme for Level III is inconsistent, though the names do match. Both authors' codes for Level III are included in the tables as a cross-reference. An earlier approach by Thom and Wilson (1980) divided Missouri into natural divisions and sections (page 7), roughly corresponding to Level III and Level IV ecoregions, respectively.

**Table 1.** List of Level I, II, and III Terrestrial Ecoregions from Omernik (1987). Those regions marked with an asterisk (\*) do not occur in Missouri but are sufficiently close to be of interest to Missouri biologists. Level III codes from Nigh and Schroeder (2002) are in parentheses.

- 8. Eastern Temperate Forests
  - 8.3. Southeastern USA Plains
    - **8.3.3.** Interior Plateau\* (71)
    - **8.3.6.** Mississippi Valley Loess Plains (74)
    - **8.3.2.** Interior River Valleys and Hills (72)
  - **8.4.** Ozark / Ouachita / Appalachian Forests
    - **8.4.5.** Ozark Highlands (39)
    - **8.4.6.** Boston Mountains\* (38)
  - 8.5. Southeast US Coastal Plain
    - **8.5.2.** Mississippi Alluvial Plain (73)

- 9. Great Plains
  - **9.2.** Temperate Prairies
    - **9.2.3.** Western Corn Belt Plains (47)
    - **9.2.4.** Central Irregular Plains (40)

**Table 2.** List of Level III and Level IV Terrestrial Ecoregions from Nigh and Schroeder (2002). Those regions marked with an asterisk (\*) do not occur in Missouri but are sufficiently close to be of interest to Missouri biologists. Those regions marked with a caret (^) are contained entirely within the state's borders.

- **38.** Boston Mountains\* (8.4.6)
  - **38a.** Upper Boston Mountains\*
  - **38b.** Lower Boston Mountains\*
- **39.** Ozark Highlands (8.4.5)
  - 39a. Springfield Plateau
  - 39b. Elk River Hills
  - **39c.** White River Hills
  - **39d.** Central Plateau
  - 39e. Osage / Gasconade River Hills^
  - **39f.** Saint François Knobs / Basins^
  - 39q. Meramec River Hills^
  - 39h. Current River Hills^
  - 39i. Easter Ozark Border^
  - **39j.** Black River Hills Border^
  - 39k. Prairie Ozark Border^
- **40.** Central Irregular Plains (9.2.4)
  - 40a. Loess Flats and Till Plains
  - 40b. Osage Cuestas\*
  - **40c.** Wooded Osage Plains
  - **40d.** Cherokee Plains
  - 40e. Claypan Prairie^
- **47.** Western Corn Belt Plains (9.2.3)
  - 47d. Missouri Alluvial Plain
  - 47e. Steeply Rolling Loess Prairies
  - 47f. Rolling Loess Prairies
  - **47h**. Nebraska / Kansas Loess Hills\*
  - 47i. Loess and Glacial Drift Hills\*
  - 47m. Western Loess Hills
- **71.** Interior Plateau\* (8.3.3)
  - **71m.** Northern Shawnee Hills\*
  - 71n. Southern Shawnee Hills\*
- **72**. Interior River Valleys and Hills (8.3.2)
  - 72a. Wabash / Ohio Bottomlands\*
  - **72d.** Upper Mississippi Alluvial Plain
  - **72e.** Middle Mississippi Alluvial Plain
  - 72f. River Hills
  - 72g. Southern Ozarkian River Bluffs\*
  - 72i. Western Dissected Illinoian Till Plain\*
  - 72 j. Southern Illinoian Till Plain\*
  - 72k. Cretaceous Hills\*
  - 721. Karstic Northern Ozarkian River Bluffs\*
- **73.** Mississippi Alluvial Plain (8.5.2)
  - **73a.** Northern Holocene Meander Belts
  - 73b. Northern Pleistocene Valley Trains\*
  - 73c. Saint Francis Lowlands
  - **73f.** Western Lowlands Holocene Meander Belts

73g. Western Lowlands Pleistocene Valley Trains

**74.** Valley Loess Plains (8.3.6)

74a. Bluff Hills74b. Loess Plains

### **Aquatic Subregions**

Like terrestrial ecoregions, aquatic subregions can be classified in discrete units, based on watershed and hydrological characteristics. Subregions are composed of ecological drainage units, which are major watersheds that are consolidated based on similarity and proximity (MSDIS 2009a). Subregions and their component drainage units can be seen in Table 3 (below).

Naturally, many of the same patterns emerge in both terrestrial and aquatic geographic classifications since the two are not independent of one another. However, aquatic subregions and ecological drainage units can differ from their terrestrial counterparts since most aquatic organisms are confined to watersheds as well as specific habitats. Since many species of amphibians and reptiles are partially or wholly dependent on aquatic habitats, distributions can sometimes be best understood and explained by examining aquatic subregions, ecological drainage units, and even individual watersheds.


The base map for major rivers and streams (page 12) shows locations of actual rivers and streams inside Missouri and just outside the state's borders where appropriate (USGS 1994; MDC 2007b). The watershed boundaries map (page 13) displays the maximum terrestrial boundary for major watersheds (NRCS 2002). Both of these maps are underlain with unique background colors to display Missouri's primary drainage systems: a) Mississippi River, b) Missouri River, c) Arkansas River, and d) White River.


Some species found in the state exhibit distributions that can be better understood by referring to these primary drainage systems. The ecological drainage units map (page 14) shows a somewhat consolidated view of the watersheds map based on those watersheds and other characteristics. It is underlain with unique background colors showing the three broad aquatic subregions in the state. Still more detailed aquatic region classifications are available but often include areas based on very small creeks and their watersheds and are therefore not included here.


**Table 3.** List of Aquatic Subregions and Ecological Drainage Units from MSDIS (2009a, 2009c). Each drainage unit is indicated with its primary drainage system: a) Mississippi River, b) Missouri River, c) Arkansas River, d) White River.


- \* Central Plains Subregion
  - \* Blackwater / Lamine (b)
  - \* Cuivre / Salt (a)
  - \* Des Moines (a)
  - \* Grand / Chariton (b)
  - \* Kansas (b)
  - \* Nishnabotna / Platte (b)
  - \* Osage / South Grand (b)
- Ozarks Subregion


- \* Apple / Joachim (a)
- \* Black / Current (d)
- \* Gasconade (b)
- \* Meramec (a)
- \* Moreau / Loutre (b)
- \* Osage (b)
- \* Neosho (c)
- \* Upper Saint Francis / Castor (a)
- \* White (d)
- \* Mississippi Alluvial Subregion
  - \* Black / Cache (a)
  - \* Saint Francis / Little (a)
  - \* Saint Johns Bayou (a)

























### **Species Locality Maps**

Key:

- Museum or published record believed to be valid
- \* County record only for:
  - -- Valid record without a specific locality
  - -- Species of conservation concern, locality withheld
- ? Questionable record that may be valid




Siren intermedia
Western Lesser Siren




Cryptobranchus alleganiensis Hellbender

Ambystoma annulatum Ringed Salamander




Ambystoma maculatum Spotted Salamander

Ambystoma opacum Marbled Salamander




Ambystoma talpoideum Mole Salamander

Ambystoma texanum Small-mouthed Salamander




Ambystoma tigrinum Eastern Tiger Salamander

Notophthalmus viridescens Central Newt




Necturus maculosus Mudpuppy

Amphiuma tridactylum
Three-toed Amphiuma




Eurycea longicauda Long-tailed Salamander

Eurycea lucifuga Cave Salamander




Eurycea spelaea Grotto Salamander

Eurycea tynerensis Oklahoma Salamander




Hemidactylium scutatum Four-toed Salamander


Plethodon albagula Western Slimy Salamander




Plethodon angusticlavius Ozark Zigzag Salamander

Plethodon serratus
Southern Red-backed Salamander










Hyla chrysoscelis Cope's Gray Treefrog Acris blanchardi

Blanchard's Cricket Frog









Pseudacris fouquettei Cajun Chorus Frog


Pseudacris illinoensis
Illinois Chorus Frog




Pseudacris maculata Boreal Chorus Frog

Lithobates areolatus Northern Crawfish Frog









Lithobates pipiens
Northern Leopard Frog

Lithobates sphenocephalus
Southern Leopard Frog




Lithobates sylvaticus Wood Frog

Gastrophryne carolinensis
Eastern Narrow-mouthed Toad




Gastrophryne olivacea Western Narrow-mouthed Toad

Crotaphytus collaris Eastern Collared Lizard




Phrynosoma cornutum Texas Horned Lizard

Sceloporus consobrinus Prairie Lizard




Aspidoscelis sexlineata Six-lined Racerunner

Plestiodon anthracinus Southern Coal Skink




Plestiodon fasciatus Common Five-lined Skink

Plestiodon laticeps Broad-headed Skink




Plestiodon obsoletus Great Plains Skink

Plestiodon septentrionalis Prairie Skink




Scincella lateralis Little Brown Skink

Ophisaurus attenuatus Western Slender Glass Lizard




Hemidactylus turcicus Mediterranean Gecko

Podarcis siculus Italian Wall Lizard




Cemophora coccinea Northern Scarletsnake

Coluber constrictor
North American Racer




Coluber flagellum Eastern Coachwhip

Lampropeltis calligaster Prairie Kingsnake




Lampropeltis holbrooki Speckled Kingsnake

Lampropeltis nigra Eastern Black Kingsnake




Lampropeltis triangulum Eastern Milksnake

Opheodrys aestivus Northern Rough Greensnake




Opheodrys vernalis Smooth Greensnake

Pantherophis emoryi Great Plains Ratsnake




Pantherophis obsoletus Western Ratsnake

Pantherophis ramspotti Western Foxsnake




Pantherophis vulpinus Eastern Foxsnake

Pituophis catenifer
Bullsnake




Sonora semiannulata Variable Groundsnake

Tantilla gracilis Flat-headed Snake




Carphophis vermis Western Wormsnake

Diadophis punctatus Ring-necked Snake




Farancia abacura Western Mudsnake

Heterodon gloydi
Dusty Hog-nosed Snake




Heterodon nasicus
Plains Hog-nosed Snake

Heterodon platirhinos Eastern Hog-nosed Snake




Clonophis kirtlandii Kirtland's Snake

Haldea striatula Rough Earthsnake




Nerodia cyclopion Mississippi Green Watersnake

Nerodia erythrogaster Plain-bellied Watersnake




Nerodia fasciata
Broad-banded Watersnake

Nerodia rhombifer
Diamond-backed Watersnake




Nerodia sipedon Common Watersnake

Regina grahamii Graham's Crawfish Snake




DeKay's Brownsnake

Storeria occipitomaculata Northern Red-bellied Snake




Thamnophis proximus
Orange-striped Ribbonsnake

Thamnophis radix
Plains Gartersnake




Thamnophis sirtalis
Common Gartersnake

Tropidoclonion lineatum Lined Snake




Virginia valeriae Western Smooth Earthsnake

Agkistrodon contortrix
Copperhead




\*\*\*

Sistrurus catenatus Eastern Massasauga Sistrurus miliarius Western Pygmy Rattlesnake




Sistrurus tergeminus Prairie Massasauga

Chelydra serpentina Snapping Turtle




Macrochelys temminckii Alligator Snapping Turtle

Kinosternon flavescens Yellow Mud Turtle




Mississippi Mud Turtle

Sternotherus odoratus Eastern Musk Turtle




Chrysemys dorsalis
Southern Painted Turtle

Chrysemys picta Western Painted Turtle




Deirochelys reticularia Western Chicken Turtle

Emydoidea blandingii Blanding's Turtle




Graptemys geographica Northern Map Turtle

Graptemys ouachitensis Ouachita Map Turtle




Graptemys pseudogeographica False Map Turtle

Pseudemys concinna River Cooter




Terrapene carolina
Three-toed Box Turtle

Terrapene ornata Ornate Box Turtle



Trachemys scripta
Red-eared Slider

Apalone mutica
Midland Smooth Softshell




Apalone spinifera
Eastern Spiny Softshell


# **County Records Maps**

Key:

Valid vouchered county record




Siren intermedia
Western Lesser Siren




Cryptobranchus alleganiensis Hellbender

Ambystoma annulatum Ringed Salamander




Ambystoma maculatum Spotted Salamander

Ambystoma opacum Marbled Salamander




Ambystoma talpoideum Mole Salamander

Ambystoma texanum Small-mouthed Salamander




Ambystoma tigrinum Eastern Tiger Salamander

Notophthalmus viridescens Central Newt




Necturus maculosus Mudpuppy

Amphiuma tridactylum
Three-toed Amphiuma




Eurycea longicauda Long-tailed Salamander

Eurycea lucifuga Cave Salamander




Eurycea spelaea Grotto Salamander

Eurycea tynerensis Oklahoma Salamander




Hemidactylium scutatum Four-toed Salamander

Plethodon albagula Western Slimy Salamander




Plethodon angusticlavius Ozark Zigzag Salamander

Plethodon serratus
Southern Red-backed Salamander




Scaphiopus holbrookii Eastern Spadefoot

Spea bombifrons
Plains Spadefoot




Anaxyrus americanus American Toad

Anaxyrus cognatus Great Plains Toad




Anaxyrus fowleri Fowler's Toad

Anaxyrus woodhousii Woodhouse's Toad




Acris blanchardi Blanchard's Cricket Frog

Hyla chrysoscelis Cope's Gray Treefrog




Hyla cinerea Green Treefrog

Hyla versicolor Gray Treefrog




Pseudacris crucifer
Spring Peeper

Pseudacris feriarum Upland Chorus Frog




Pseudacris fouquettei Cajun Chorus Frog

Pseudacris illinoensis
Illinois Chorus Frog




Pseudacris maculata Boreal Chorus Frog

Lithobates areolatus Northern Crawfish Frog




Lithobates blairi
Plains Leopard Frog

Lithobates catesbeianus
American Bullfrog




Lithobates clamitans Green Frog

Lithobates palustris
Pickerel Frog




Lithobates pipiens
Northern Leopard Frog

Lithobates sphenocephalus
Southern Leopard Frog




Lithobates sylvaticus Wood Frog

Gastrophryne carolinensis
Eastern Narrow-mouthed Toad




Gastrophryne olivacea Western Narrow-mouthed Toad

Crotaphytus collaris Eastern Collared Lizard




Phrynosoma cornutum Texas Horned Lizard

Sceloporus consobrinus Prairie Lizard




Aspidoscelis sexlineata Six-lined Racerunner

Plestiodon anthracinus Southern Coal Skink




Plestiodon fasciatus Common Five-lined Skink

Plestiodon laticeps Broad-headed Skink




Plestiodon obsoletus Great Plains Skink

Plestiodon septentrionalis Prairie Skink




Scincella lateralis Little Brown Skink

Ophisaurus attenuatus Western Slender Glass Lizard




Hemidactylus turcicus Mediterranean Gecko

Podarcis siculus Italian Wall Lizard




Cemophora coccinea Northern Scarletsnake

Coluber constrictor
North American Racer




Coluber flagellum Eastern Coachwhip

Lampropeltis calligaster Prairie Kingsnake




Lampropeltis holbrooki Speckled Kingsnake

Lampropeltis nigra Eastern Black Kingsnake




Lampropeltis triangulum Eastern Milksnake

Opheodrys aestivus Northern Rough Greensnake




Opheodrys vernalis Smooth Greensnake

Pantherophis emoryi Great Plains Ratsnake




Pantherophis obsoletus Western Ratsnake

Pantherophis ramspotti Western Foxsnake




Pantherophis vulpinus Eastern Foxsnake

Pituophis catenifer Bullsnake




Sonora semiannulata Variable Groundsnake

Tantilla gracilis
Flat-headed Snake




Carphophis vermis Western Wormsnake

Diadophis punctatus Ring-necked Snake




Farancia abacura Western Mudsnake

Heterodon gloydi
Dusty Hog-nosed Snake




Heterodon nasicus
Plains Hog-nosed Snake

Heterodon platirhinos Eastern Hog-nosed Snake




Clonophis kirtlandii Kirtland's Snake

Haldea striatula Rough Earthsnake




Nerodia cyclopion Mississippi Green Watersnake

Nerodia erythrogaster Plain-bellied Watersnake




Nerodia fasciata
Broad-banded Watersnake

Nerodia rhombifer
Diamond-backed Watersnake




Nerodia sipedon Common Watersnake

Regina grahamii Graham's Crawfish Snake




Storeria dekayi DeKay's Brownsnake

Storeria occipitomaculata Northern Red-bellied Snake




Thamnophis proximus
Orange-striped Ribbonsnake

Thamnophis radix
Plains Gartersnake




Thamnophis sirtalis
Common Gartersnake

Tropidoclonion lineatum Lined Snake




Virginia valeriae Western Smooth Earthsnake

Agkistrodon contortrix
Copperhead




Agkistrodon piscivorus Western Cottonmouth

Crotalus horridus
Timber Rattlesnake




Sistrurus catenatus Eastern Massasauga

Sistrurus miliarius Western Pygmy Rattlesnake




Sistrurus tergeminus Prairie Massasauga

Chelydra serpentina Snapping Turtle




Macrochelys temminckii Alligator Snapping Turtle

Kinosternon flavescens Yellow Mud Turtle




Kinosternon subrubrum Mississippi Mud Turtle

Sternotherus odoratus Eastern Musk Turtle




Chrysemys dorsalis
Southern Painted Turtle

Chrysemys picta Western Painted Turtle




Deirochelys reticularia Western Chicken Turtle

Emydoidea blandingii Blanding's Turtle




Graptemys geographica Northern Map Turtle

Graptemys ouachitensis Ouachita Map Turtle




Graptemys pseudogeographica False Map Turtle

Pseudemys concinna River Cooter



Terrapene carolina
Three-toed Box Turtle

Terrapene ornata Ornate Box Turtle



Trachemys scripta
Red-eared Slider

Apalone mutica
Midland Smooth Softshell



Apalone spinifera
Eastern Spiny Softshell

## Acknowledgements

Data for the base county maps for the locality and county distributions, the labeled county map, and the ecological drainage units and aquatic subregions were generously provided by the Missouri Spatial Data Information Service (MSDIS 2009a, 2009b, 2009c). Data for the major rivers were provided by the United States Geological Survey (USGS 1994) and supplemented with data from the Missouri Department of Conservation (MDC 2007). Data for Missouri natural divisions and sections (based on Thom and Wilson (1980)) were generously provided by the Missouri Department of Conservation (MDC 2007). Data for watershed boundaries were provided by the Natural Resources Conservation Service (NRCS 2002). Data for ecoregions were provided by the US Environmental Protection Agency (EPA 2003). Base map data for surrounding states were provided by the National Atlas of the United States (National Atlas 2001).

All technology-related aspects of the Missouri Herpetological Atlas Project are supported by a suite of open source software that is both *gratis* (free as in beer) and *libre* (free as in speech). Kubuntu Linux is used as the operating system to host and execute all processes related to the project. Atlas, checklist, and GIS data are stored in a PostgreSQL database, which has been spatially enabled using PostGIS. Project documentation and database scripts are written with PHP and displayed using an Apache web server. QGIS is used to produce all base maps, distribution, county records, and other maps. The programming language Python is used to create GIS plug-ins for map exports. The open source Python library, ReportLab, is used to create the final *Atlas* publication in portable document format (PDF), an ISO standard.

Most importantly, a project of this magnitude cannot be accomplished without the contributions of many individuals. We would like to thank the following for specimen information, feedback, and technical assistance on the project:

Nathan Aaron, Jamie Alford, Scott Altnether, Tom Anderson, Richard Andrews, Jonathan Ashley, Ralph Axtell, D. Baker, Jason Baker, Stephanie Bartlett, Ben Barton, Neil Bass, Derek Bateman, Tom Beauvais, Bill Becker, Don Becker, Melissa Bedford, Ruth Ann Berry, Adam Beston, Beth Blaisdell, Tyler Blake, Josiah Boothe, Aaron Bossert, Michelle Bowe, Mark Bower, Jeff Bowman, Andrew Braun, Jeff Briggler, Chris Brooks, Christina Brown, Danny Brown, Sara Bruce, Josh Bruegge, Kevin Brunke, Carol Bryant, Steve Buback, Jenny Bunch, Frank Burbrink, Nick Burrell, John Cadle, T. Carpenter, Gary Casper, Robert Caviness, Ellen Censky, Candy Chambers, Doug Clark, Clint Coffey, Shayla Coffey, Skyler Collins, Matt Coombs, Wayne Corse, Danny Counts, Christopher Crabtree, Mike Cravens, Joe Cridlebaugh, Wyatt Cridlebaugh, Mary Crowell, R. Dames, Charles Dardin, Jeffrey Dawson, B. Day, Bob Dealey, Brandon DeCavele, Jacob Decker, Michael Dickey, Angel DiMercurio, Drew Dittmer, Brandon Dix, Evan Dixon, Scott Doherty, Ryan Douglas, Joseph Douglass, D. Drake, Frank Drummond, Joe Duenwald, Spencer Duffendack,

Frank Durbian, Eric East, David Easterla, Amber Edwards, T. Enright, Rick Essner, Pete Eyheralde, Susan Farrington, E. Field, Amanda Fisher, Z. Fratto, G. Frazier, Mark Frazier, Dillon Freiburger, Paul Frese, Darrell Frost, Bonnie Gardner, C. Gemming, Carl Gerhardt, Matt Gifford, Mark Gihring, Brian Gilbert, Bob Gillespie, T. Girard, Brad Glorioso, Steve Gotte, Millie Graham, Brian Greene, Greg Gremaud, Adam Guerich, Brad Hadley, Adam Hamlin, D. Harrison, Kevin Hedgpeth, AJ Hendershott, Robert Henderson, Mary Hennen, John Hess, Joshua Hill, Dan Hocking, Colby Hodges, Dana Hoisington, Ethan Hollender, Terry Holtsclaw, E. Hooper, Emily Horner, Ann Horsman, Brian Hubbs, Jamie Hubert, James Hunt, Seth Ince, Adam Ivey, Thomas Jay, Matt Jeppson, Linda Jepson, Tom Jessen, Tom Johnson, Joe Jones, Mike Jones, Kayl Kapral, John Karel, Tim Kavan, Kate Kelly, Robert Kipfer, William Kirby, Arnold Kluge, Kent Korthas, Robert Krager, Jim Lamer, Alek Lantar, Mike Leahy, Joseph Lemen, Kevin Lewis, Stephanie Liebi, Day Ligon, Elicia Ligon, Kenton Lohraff, Alexandria London, Peter Lowther, Beth Lynch, Jennifer Maag, Glenn Manning, Nicholas March, Dexter Mardis, Kevin Martie, Gina Martin, Terrell Martin, Frank Martinez, Rebecca Maxfield, Christine Mayer, Amy McClure, Roy McDiarmid, Elizabeth McGhee, B. McKeage, Paul McKenzie, Don McKnight, Arianne Messerman, Dean Metter, C. Meyer, Brian Miller, Danielle Miller, J. Miller, N. Mitchell, Chad Montgomery, Nathan Montgomery, D. Moore, T. Moore, L. Moree, Denise Morgan-Irish, Nathanael Morris, K. Morrow, Sarah Moss, Nathan Muenks, Tom Nagel, Kevin Narum, L. Neuman-Lee, Chris Newbold, Abbigail Nicholson, Krista Noel, Laurel Norton, Jim Oehmke, Tyler Olcott, Matt Ormsby, Seth Otto, Cheryl Palmer, Peter Paplanus, John Par, Ben Parnell, Caleb Pemberton, Bill Peterman, M. Peterson, Katie Petrie, Tim Pettinelli, Chris Phillips, Nick Pieroni, Gregory Pinson, G. Pitchford, P. Pitts, Jon Pollock, Crystal Poole, Robert Powell, Tracey Prenger, Yarrow Price, B. Probst, Keith Pulles, Holly Quinn, Hugh Quinn, Corey Raimond, Alan Resetar, Joe Ridings, Daren Riedle, Rhonda Rimer, Tony Robyn, Bernard Ross, Douglas Rossman, David Rostollan, Chad Runco, B. Russell, Greg Schneider, Tim Schoppenhorst, E. Schott, Bruce Schuette, Jeremy Schumacher, C. Scott, K. Scott, John Severson, Alex Sharp, Chris Shelton, Chris Shulse, Dustin Siegel, Carri Silman, John Simmons, Rachel Simmons, Carl Smith, Geoff Smith, Isaac Smith, Madeline Smith, S. Smith, Steven Snow, Leon Spring, Marianne Stanley, Clarissa Starbuck, Andy Starostka, Shane Staten, Brandon Stehl, Jeff Stewart, Greg Stoner, James Studdard, Kelly Sulkowski, Mercedies Summers, Becky Swearingen, Travis Taggart, Jeremiah Terrell, Wendell Tharp, Ryan Thies, M. Thompson, A. Thorn, Bart Tibbs, Mark Tovar, Sean Tracy, Stan Trauth, Angelina Trombley, T. Trombone, Linda Trueb, Joe Twardowski, Chad Wallis, Sammy Wentz, J. West, Steve Westin, Amber Wiewel, Brian Willey, Bethany Williams, Linda Williams, Myki Williamson, Matt Wishard, Rod Wittenberg, N. Woodland, Nathan Woolery, Reese Worthington, Patricia Yates, Nico Yoch, Elizabeth Yohe

#### Literature Cited

- Anderson, P. 1945. New herpetological records for Missouri. Bulletin of Chicago Academy of Sciences 7(5): 271-275.
- Anderson, P. 1957. A second list of new herpetological records for Missouri. Natural History Miscellanea (161): 1-5.
- Anderson, P. 1965. *The Reptiles of Missouri*. University of Missouri Press, Columbia, Missouri. xxiii + 330p.
- Briggler, J.T. and T.R. Johnson. 2021. *The Amphibians and Reptiles of Missouri* (3rd ed.). Missouri Department of Conservation, Jefferson City, Missouri.
- Briggler, J.T., R.L. Rimer and G. Deichsel. 2015. First record of the Northern Italian Wall Lizard (*Podarcis siculus campestris*) in Missouri. *IRCF Reptiles and Amphibians* 22(1): 43-45.
- Bufalino, A.P. 2004. Geographic Distribution: *Hemidactylus turcicus turcicus*. *Herpetological Review* 35: 188.
- Burke, R.L. and G. Deichsel. 2008. Lacertid lizards introduced into North America: History and future, pp.347-353. In: J.C. Mitchell, R.E.I. Brown and B. Bartholomew (eds.) *Urban Herpetology*. Herpetological Conservation vol. 3. SSAR, Salt Lake City, Utah.
- Conant, R. 1960. The Queen Snake, Natrix septemvittata, in the Interior Highlands of Arkansas and Missouri, with comments upon similar disjunct distributions. Proceedings of The Academy of Natural Sciences of Philadelphia 112(2): 25-40.
- Conant, R., and J.T. Collins. 1998. A Field Guide to Reptiles and Amphibians: Eastern and Central North America. (3rd ed., expanded) Peterson Field Guides, Houghton Mifflin Company, New York. 616p.
- Daniel, R.E., and B.S. Edmond. 2002. Revised County Distribution Maps of Amphibians and Reptiles of Missouri. Missouri Herpetological Association Newsletter 15: 16-38.
- Daniel, R.E., and B.S. Edmond. 2022. Atlas of Missouri Amphibians and Reptiles for 2021. <a href="https://atlas.moherp.org/pubs/atlas21.pdf">https://atlas.moherp.org/pubs/atlas21.pdf</a>
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2015. New herpetological distribution records for Missouri in 2015. Missouri Herpetological Association Newsletter 28: 10-14.
- Daniel, R.E., B.S. Edmond, and J.T. Briggler. 2022. New herpetological distribution records for Missouri in 2022. *Missouri Herpetological Association Newsletter* 35: 3-5.
- Edmond, B.S. and R.E. Daniel. 2024.
  - Missouri Herpetological Atlas Project. <a href="http://atlas.moherp.org/">http://atlas.moherp.org/</a> Accessed 19 April 2024.
- Environmental Protection Agency (EPA). 2003. Level III and IV Ecoregions of the Continental United States. <a href="http://www.epa.gov/wed/pages/ecoregions/na\_eco.htm">http://www.epa.gov/wed/pages/ecoregions/na\_eco.htm</a> Accessed 1 December 2009.
- Hawker, J.L. 1992. Missouri Landscapes: A Tour Through Time. Missouri Department of Natural Resources, Division of Geology and Land Survey, Rolla. 326pp.
- Hurter, J. 1911. Herpetology of Missouri. *Transactions of the Saint Louis Academy of Science* 20(5): 59-274.
- Johnson, T.R. 1987. The Amphibians and Reptiles of Missouri. (1st ed.) Missouri Department of Conservation, Jefferson City, Missouri. 368p.
- Johnson, T.R. 2000. The Amphibians and Reptiles of Missouri. (2nd ed.) Missouri Department of Conservation, Jefferson City, Missouri. 400p.
- Johnson, T.R. and R.N. Bader. 1974. Annotated checklist of Missouri amphibians and reptiles. *St. Louis Herpetological*

- Society Special Issue 1: 1-16.
- Missouri Department of Conservation (MDC). 2007a. *Natural Divisions and Sections*. <a href="http://www.mdc.mo.gov/">http://www.mdc.mo.gov/</a> Accessed 1 February 2007.
- Missouri Department of Conservation (MDC). 2007b. Rivers and Streams. <a href="http://www.mdc.mo.gov/">http://www.mdc.mo.gov/</a> Accessed 1 February 2007.
- Missouri Spatial Data Information Service (MSDIS). 2009a. Aquatic Subregions <a href="http://msdis.missouri.edu/">http://msdis.missouri.edu/</a> Accessed 1 December 2009.
- Missouri Spatial Data Information Service (MSDIS). 2009b. *Base County and Boundary Map* <a href="http://msdis.missouri.edu/">http://msdis.missouri.edu/</a> Accessed 1 December 2009.
- Missouri Spatial Data Information Service (MSDIS). 2009c. *Ecological Drainage Units* <a href="http://msdis.missouri.edu/">http://msdis.missouri.edu/</a> Accessed 1 December 2009.
- National Atlas of the United States. 2001. State and County Boundaries of the United States. <a href="http://nationalatlas.gov/atlasftp.html">http://nationalatlas.gov/atlasftp.html</a> Accessed 1 December 2009.
- Natural Resources Conservation Service, U.S. Department of Agriculture. 2002. Watershed Boundaries of the United States. <a href="http://datagateway.nrcs.usda.gov/">http://datagateway.nrcs.usda.gov/</a> Accessed 1 December 2009
- Nelson, P.W. 2005. *The Terrestrial Natural Communities of Missouri*. Missouri Natural Areas Committee, Jefferson City. 550pp.
- Nickerson, M.A. and R. Krager. 1972. Additional noteworthy records of Missouri amphibians and reptiles with a possible addition to the herpetofauna. *Transactions of the Kansas Academy of Science* 75(3) 276-277.
- Nigh, T.A., and W.A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City. 212pp.
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. Map Supplement (scale 1:7,500,000). *Annals of the Association of American Geographers* 77(1): 118-125.
- Pflieger, W.L. 1989. Aquatic community classification system for Missouri. *Aquatic Series No. 19*. Missouri Department of Conservation, Jefferson City.
- Powell, R., R. Conant, and J.T. Collins. 2016. *Peterson Field Guide to Reptiles and Amphibians of Eastern and Central North America*. (4th ed.) Houghton Mifflin Harcourt Publishing Company, New York, 494p.
- Powell, R. and R. Daniel. 1997. County distribution records of amphibians and reptiles in Missouri. *Missouri Herpetological Association Newsletter* 10: 12-69.
- Powell, R., T.R. Johnson, and J.S. Parmerlee. 1993. Updated distribution maps for amphibians and reptiles in *Missouri Herpetological Association Newsletter* 6: 7-26.
- Shepard, D.B. and A.R. Kuhns. 2017. Shifting rivers and stationary ground: biogeographic history of slimy salamanders on an isolated bluff in the Mississippi River floodplain. *Journal of Zoology* DOI:10.1111/jzo.12514
- Steyermark, J.A. 1963. *Flora of Missouri*. Iowa State University Press, Ames. 1728pp.
- Thom, R.H., and J.H. Wilson. 1980. The natural divisions of Missouri. *Transactions of the Missouri Academy of Science* 14:9-23.
- United States Geological Survey. 1994. Major Rivers and Streams of the United States. <a href="http://pubs.usgs.gov/">http://pubs.usgs.gov/</a> Accessed 1

December 2009.

Unklesbay, A.G., and J.D. Vineyard. 1992. *Missouri Geology: Three Billion Years of Volcanoes, Seas, Sediments, and Erosion*. University of Missouri Press, Columbia. 189pp.
Yatskievych, G. 1999. *Steyermark's Flora of Missouri, Volume 1*.

Missouri Department of Conservation, Jefferson City. 991pp.

### **Appendix A: Collection Sources**

American Museum of Natural History

Arkansas State University

Austin Peay State University \*

Bobby Witcher Memorial Collection, Avila College (KU)

Carnegie Museum of Natural History

Chicago Academy of Sciences

Cornell University Museum of Vertebrates

Culver-Stockton College

Dallas Museum of Natural History \*

Field Museum of Natural History

H. Carl Gerhardt Audio Recording Collection (Cornell)

Illinois Natural History Survey

Milwaukee Public Museum

Missouri Department of Conservation \*

Missouri State University

Museum of Comparative Zoology, Harvard University

Museum of Natural Science, Louisiana State University

Museum of Southwest Biology (New Mexico) \*

National Museum of Natural History

Natural History Museum of Los Angeles County \*

Pittsburg State University

Saint Louis Academy of Sciences

Southeast Missouri State University

Southern Illinois University--Carbondale

Southern Illinois University--Edwardsville

Sternberg Museum of Natural History, Fort Hays State University

Texas Natural History Collection \*

University of Arkansas--Fayetteville

University of California--Berkeley Museum of Vertebrate Zoology

University of Central Missouri

University of Florida at Gainesville \*

University of Georgia, Museum of Natural History

University of Illinois Natural History Museum

University of Kansas, Museum of Natural History

University of Michigan, Museum of Zoology

University of Missouri--Columbia Dean E. Metter Memorial Herpetology Collection

William Jewell College

<sup>\*</sup> Collection sources marked with an asterisk (\*) are included from published accounts only and are presumed to represent only part of the collection's Missouri records. All Missouri material from these sources has not been included in the database or the printed atlas.

### **Appendix B: Literature Sources**

Anderson, P. 1942. Amphibians and reptiles of Jackson County, Missouri. *Bulletin of the Chicago Academy of Science* 6(11): 203-220.

Anderson, P. 1965. *The Reptiles of Missouri*. University of Missouri Press, Columbia, Missouri. xxiii + 330p.

Blanchard, F.N. 1924. A collection of amphibians and reptiles from southeastern Missouri and Illinois. *Papers of the Michigan Academy of Science, Art, and Letters* 4: 533-541.

Bodie, J.R. 1998. Contemporary hydrology of a large-river floodplain: Multiple-scale effects on freshwater turtles. Unpublished MA thesis, University of Missouri-Columbia.

Boyer, D.A., and A.A. Heinze. 1934. An annotated list of the amphibians and reptiles of Jefferson County, Missouri. *Transactions of the Saint Louis Academy of Science* 28(4): 185-200.

Bufalino, A.P. 2004. Geographic Distribution: *Hemidactylus turcicus turcicus*. *Herpetological Review* 35: 188.

Burt, C.E. 1932. Records of amphibians from the eastern and central United States (1931). *American Midland Naturalist* 13: 75-85.

Burt, C.E. 1933. A collection of amphibians and reptiles from southern Missouri. *American Midland Naturalist* 14: 170-173.

Burt, C.E. 1935. Further records of the ecology and distribution of amphibians and reptiles in the middle west. *American Midland Naturalist* 16(3): 311-366.

Burt, C.E. and W.L. Hoyle. 1935. Additional records of the reptiles of the central prairie region of the United States. *Transactions of the Kansas Academy of Science* 37: 193-216.

Conant, R., and C.J. Goin. 1948. A new subspecies of the soft-shelled turtle from the central United States, with comments on the application of the name *Amyda*. *Occasional Papers of the Museum of Zoology* (510): 1-19.

Easterla, D.A. 1972. Herpetological records for northwest Missouri. *Transactions of the Missouri Academy of Science* 6: 158-160.

Easterla, D.A. and M.D. Meadows. 1993. The northern prairie skink rediscovered in Missouri, the last recorded occurrence of the smooth green snake for Missouri, plus a Missouri range extension for the northern spring peeper. *Bluebird* 60(3): 105-109.

Evans, P.D. 1940. Notes on Missouri snakes. *Copeia* 1940: 53-54

Henning, W.L. 1938. Amphibians and reptiles of a 2,220-acre tract in central Missouri. *Copeia* 1938(2): 91-92.

Hubbs, B. 2011. Geographic Distribution: *Lampropeltis triangulum syspila*. *Herpetological Review* 42: 114.

Hubbs, B. 2012. Geographic Distribution: *Apalone mutica*. *Herpetological Review* 43(3): 441.

Hurter, J. 1911. Herpetology of Missouri. *Transactions of the Saint Louis Academy of Science* 20(5): 59-274.

Jones, J.M. 1967. A western extension of the known range of Kirtland's watersnake. *Herpetologica* 23(1): 66-67.

Karel, J.A. 1974. Cottonmouths in the St. Francois Mountains. *Saint Louis Herpetological Society Newsletter* 1(8): 10.

McConkey, E.H. 1954. A systematic study of the North American lizards of the genus *Ophisaurus*. *American Midland Naturalist* 51(1): 133-171.

Myers, C.W. 1957. Amphibians and reptiles of Washington State Park, Washington County, Missouri. *Transactions of the Kansas Academy of Science* 60(3): 288-293.

Myers, C.W. 1959. Amphibians and reptiles of Montauk State Park and vicinity, Dent County, Missouri. *Transactions of the Kansas Academy of Science* 62(1): 88-90.

Nicholson, A.E. 2013a. Geographic distribution: USA: Missouri: *Notophthalmus viridescens. Herpetological Review* 44(3): 470.

Nicholson, A.E. 2013b. Geographic distribution: USA: Missouri: *Plestiodon laticeps. Herpetological Review* 44(3): 475.

Nicholson, A.E. 2013c. Geographic distribution: USA: Missouri: *Lampropeltis triangulum. Herpetological Review* 44(3): 477.

Nickerson, M.A., and R. Krager. 1971. Noteworthy records of Missouri reptiles. *Transactions of the Kansas Academy of Science* 74(1): 99-101.

Nickerson, M.A. and R. Krager. 1972. Additional noteworthy records of Missouri amphibians and reptiles with a possible addition to the herpetofauna. *Transactions of the Kansas Academy of Science* 75(3) 276-277.

Owens, V. 1949. New snake records and notes, Morgan County, Missouri. *Herpetologica* 5: 49-50.

Peters, J.A. 1946. Records of certain North American salamanders. *Copeia* 1946(2): 106.

Peters, J.A. 1946. Reptiles and amphibians of Sam A. Baker State Park, Wayne County, Missouri. *Copeia* 1946(1): 44.

Phillips, C.A. 1994. Geographic distribution of mitochondrial DNA variants and the historical biogeography of the spotted salamander, *Ambystoma maculatum*. *Evolution* 48(3): 597-607.

Shaffer, H.B., and M.L. McKnight. 1996. The polytypic species revisited: Genetic differentiation and molecular phylogenetics of the tiger salamander *Ambystoma tigrinum* (Amphibia: Caudata) complex. *Evolution* 50(1): 417-433.

Tanner, W.W., and R.B. Loomis. 1957. Taxonomic and distributional study of the western subspecies of the milksnake, *Lampropeltis doliata*. *Transactions of the Kansas Academy of Science* 60(1): 12-42.

Van Wagenen, G. 1917. *Diadophis amabilis* in Missouri. *Copeia* (43): 43-44.

Watkins, L.C. 1969. A third record of the four-toed salamander, *Hemidactylium scutatum*, in Missouri. *Transactions of the Kansas Academy of Science* 72(2): 264-265.

Watkins, L.C., and L.L.Hinesley. 1970. Notes on the distribution and abundance of the sonoran skink, *Eumeces obsoletus*, in western Missouri. *Transactions of the Kansas Academy of Science* 73(1): 118-119.

Williams, K.L. 1978. American Milk Snake. *Milwaukee Public Museum Publications in Biology and Geology* 2: 1-258.